


DEC 10 2002

0611-L-02

I 49.77/5: B 87/CHAP. 14/DRAFT



U.S. Fish & Wildlife Service



Bull Trout

Salvelinus confluentus

Draft Recovery Plan

Chapter 14

Malheur River Recovery Unit

2003-009293/NF

✓

Chapter: 14

State(s): Oregon

Recovery Unit Name: Malheur Recovery Unit

**Region 1
U.S. Fish and Wildlife Service
Portland, Oregon**

I

DISCLAIMER

Recovery plans delineate reasonable actions that are believed necessary to recover and protect listed species. Plans are prepared by the U.S. Fish and Wildlife Service, and in this case, with the assistance of recovery unit teams, State and Tribal agencies, and others. Objectives will be attained and any necessary funds made available subject to budgetary and other constraints affecting the parties involved, as well as the need to address other priorities. Recovery plans do not necessarily represent the views nor the official positions or approval of any individuals or agencies involved in plan formulation, other than the U.S. Fish and Wildlife Service. Recovery plans represent the official position of the U.S. Fish and Wildlife Service only after they have been signed by the Director or Regional Director as approved. Approved recovery plans are subject to modification as dictated by new findings, changes in species status, and the completion of recovery tasks.

Literature Citation: U.S. Fish and Wildlife Service. 2002. Chapter 14, Malheur Recovery Unit, Oregon. 71 p. In: U.S. Fish and Wildlife Service. Bull Trout (*Salvelinus confluentus*) Draft Recovery Plan. Portland, Oregon.

ACKNOWLEDGMENTS

Members of the Malheur Recovery Unit Team who assisted in the preparation of this chapter include:

Wayne Bowers, Oregon Department of Fish and Wildlife
Jason Fenton, Burns Paiute Tribe
Tom Friedrichsen, U. S. Forest Service, Burns Ranger District
Dan Gonzalez, Burns Paiute Tribe
Gina Lampman, Bureau of Land Management
Sam Lohr, U.S. Fish and Wildlife Service
Alan Mauer, U.S. Fish and Wildlife Service
Steve Namitz, Burns Paiute Tribe
Gina Lampman, U. S. Bureau of Land Management, Burns District
Ray Perkins, Oregon Department of Fish and Wildlife
Rick Rieber, U.S. Bureau of Reclamation
Herb Roerick, U.S. Forest Service, Prairie City Ranger District
Tammy Salow, U.S. Bureau of Reclamation
Lawrence Schwabe, Burns Paiute Tribe
Del Skeesick, Private Citizen
Jim Soupir, U.S. Forest Service
Cynthia Tait, U.S. Bureau of Land Management, Vale District
Rick Vetter, U.S. Forest Service, Burns Ranger District
Cindy Weston, U. S. Bureau of Land Management, Burns District

Additional review and comments were provided by:

Kathryn Kostow, Oregon Department of Fish and Wildlife

MALHEUR RECOVERY UNIT

CHAPTER OF THE BULL TROUT RECOVERY PLAN

EXECUTIVE SUMMARY

CURRENT SPECIES STATUS

The U.S. Fish and Wildlife Service issued a final rule listing the Columbia River and Klamath River populations of bull trout (*Salvelinus confluentus*) as a threatened species under the Endangered Species Act on June 10, 1998 (63 FR 31647).

The Malheur Recovery Unit encompasses the entire Malheur River Basin. The basin is situated in eastern Oregon bordered on the south by the Owyhee River Basin, on the north by the Burnt River and John Day River basins, on the west by the Malheur Lakes Basin, and by the Snake River to the east, which it enters near Ontario, Oregon. The Malheur River drains an area of approximately 11,940 square kilometers (5,000 square miles) (Malheur-Owyhee Watershed Council 1999), and is approximately 306 kilometers (190 miles) from its headwaters to the confluence with Snake River (Malheur-Owyhee Watershed Council 1999).

For purposes of recovery, the Malheur Recovery Unit contains one core area, the Malheur Core Area, which includes two local populations located in the headwaters of the North Fork Malheur River and the Upper Malheur River subbasins, and the mainstem Malheur River from headwaters downstream to Namorf Dam, respectively.

HABITAT REQUIREMENTS AND LIMITING FACTORS

A detailed discussion of bull trout biology and habitat requirements is provided in Chapter 1 of this recovery plan. The limiting factors discussed here are specific to the Malheur Recovery Unit chapter. Within the Malheur Recovery Unit, historical and current land use activities have impacted bull trout local populations. There have been a combination of human-induced factors that affect bull trout including dams, forest management practices, irrigation withdrawals, livestock grazing, past bull trout harvest, and introduction of nonnative species. Lasting effects from some, but not all, of these activities still act to limit bull trout production in both local populations.

RECOVERY GOAL AND OBJECTIVES

The goal for recovery of bull trout is to **ensure the long-term persistence of self-sustaining, complex interacting groups of bull trout distributed throughout the species' native range so that the species can be delisted.** To achieve this goal the following objectives have been identified for bull trout in the Malheur River Recovery Unit:

- **Maintain the current distribution of bull trout within the core area and re-establish bull trout in previously occupied habitats in the Upper Malheur River and tributaries and the North Fork Malheur River and tributaries.**
- **Maintain stable or increasing trends in abundance of bull trout in the Malheur Recovery Unit. This will require increasing abundance within the two local populations (Upper Malheur and North Fork Malheur).**
- **Restore and maintain suitable habitat conditions for all bull trout life history stages and strategies.**
- **Conserve genetically diverse populations of bull trout populations within the Malheur Recovery Unit. This can best be achieved by ensuring connectivity between the North Fork Malheur River and the Upper Malheur River.**

RECOVERY CRITERIA

Recovery criteria for the Malheur Recovery Unit reflect the stated objectives, evaluation of population status, and recovery actions necessary to achieve the overall goal. Recovery criteria identified for the Malheur Recovery Unit are as follows:

1. **Bull trout are distributed among 2 or more local populations in the recovery unit within the Malheur Core Area. In a recovered condition one or more local populations would occur within the Upper Malheur River and one or more local populations would occur within the North Fork Malheur River.**
2. **Estimated abundance of adult bull trout in the Malheur Recovery Unit is between 2,000 and 3,000 individuals distributed between the two local populations. The recovery unit team expects to achieve this criteria by securing the current population and increasing the abundance of migratory adults in Upper Malheur River. Recovered abundance range was derived using the professional judgement of the Malheur Recovery**

Unit Team and estimation of productive capacity of identified local populations. These goals may be refined as more information becomes available, through monitoring and research.

3. **Adult bull trout exhibit a stable or increasing trend for at least 2 generations at or above the recovered abundance level within the Malheur Core Area.** The development of a standardized monitoring and evaluation program which would accurately describe trends in bull trout abundance is identified as a priority research need. As part of the overall recovery effort, the U.S. Fish and Wildlife Service will take the lead in addressing this research need by forming a multi-agency technical team to develop protocols to evaluate trends in bull trout populations.
4. **Specific barriers inhibiting bull trout movement and recovery in the Malheur Recovery Unit have been addressed, ensuring opportunities for connectivity among local populations within the core area.** Fish passage needs to be provided at Agency Dam on the North Fork Malheur and Warm Springs Dam on the Malheur River. Additional studies will be needed to determine the feasibility of providing two-way passage at Beulah and Warm Springs Dams. Reduction or elimination of threats from brook trout interaction in the Upper Malheur River will need to be accomplished prior to restoration of passage to ensure the success of bull trout recovery.

Recovery criteria for the Malheur Recovery Unit were established to assess whether recovery actions have resulted in the recovery of bull trout. Recovery criteria developed for bull trout address quantitative measurements of bull trout distribution and population characteristics. The recovery objectives were based on our current knowledge and may be refined as more information becomes available. While removal of bull trout as a species under the Endangered Species Act (*i.e.*, delisting) can only occur for the entity that was listed (Columbia River Distinct Population Segment), the recovery unit criteria listed above will be used to determine when the Malheur Recovery Unit is fully contributing to recovery of the population segment.

ACTIONS NEEDED

Recovery for bull trout will entail reducing threats to the long-term persistence of populations and their habitats, ensuring the security of multiple interacting groups of bull trout, and providing habitat and access to conditions that allow for the expression of various life history forms. Seven categories of actions needed are discussed in Chapter 1; tasks specific to this recovery unit are provided in this chapter.

ESTIMATED COST OF RECOVERY

Total estimated cost of bull trout recovery in the Malheur Recovery Unit is estimated at approximately \$10 million spread over a 25 year recovery period. Successful recovery of bull trout in the recovery unit is contingent on reconnecting bull trout populations within the recovery unit, improving habitat conditions, and removing threats from brook trout interaction and hybridization within the Upper Malheur River and tributaries. These costs are attributed to bull trout conservation, but other aquatic species will also benefit.

ESTIMATED DATE OF RECOVERY

Expected times necessary to achieve recovery will vary among recovery units due to differences in bull trout status, factors affecting bull trout, implementation and effectiveness of recovery tasks, and responses to recovery tasks. Three to five bull trout generations (15 to 25 years), or possibly longer, may be necessary before identified threats to the species can be significantly reduced and bull trout can be considered eligible for delisting. In the Malheur Recovery Unit, bull trout currently exist in very low numbers in some local populations, degradation and fragmentation of bull trout habitat presents significant migratory challenges for fluvial fish, and hybridization with introduced brook trout is occurring. Ultimately, these threats must be addressed for recovery to be achieved.

TABLE OF CONTENTS

DISCLAIMER	ii
ACKNOWLEDGMENTS	iii
EXECUTIVE SUMMARY	iv
TABLE OF CONTENTS	viii
INTRODUCTION	1
Recovery Unit Designation	1
Geographic Description	1
DISTRIBUTION AND ABUNDANCE	6
Status of Bull Trout at the Time of Listing	6
Current Distribution and Abundance	6
REASONS FOR BULL TROUT DECLINE	11
Dams	12
Forest Management Practices	14
Livestock Grazing	17
Agricultural Practices	18
Transportation Network	19
Mining	20
Residential Development	20
Fisheries Management	20
Isolation and Habitat Fragmentation	22
ONGOING RECOVERY UNIT CONSERVATION MEASURES	23
RELATIONSHIP TO OTHER CONSERVATION EFFORTS	27
State of Oregon	27
Burns Paiute Tribe	28
Local Planning Efforts	28
Northwest Power Planning Council's Subbasin Planning	29
STRATEGY FOR RECOVERY	30
Recovery Goals and Objectives	31
Recovery Criteria	35
Research Needs	36
ACTIONS NEEDED	39
Recovery Measures Narrative	39

IMPLEMENTATION	49
REFERENCES	58
APPENDIX A: Historical and current fish species found in the Malheur River Basin excerpted from NWPPC 2001	63
APPENDIX B: List of barriers in the Malheur Recovery Unit known or suspected to impact bull trout.	65
APPENDIX C: List of Instream Water Rights established to benefit bull trout (BuT) and redband trout (RT).	66
APPENDIX D: Oregon Department of Fish and Wildlife instream water right priorities for bull trout (BuT) and redband trout (RT).	67
APPENDIX E: List of current and potential (*) local bull trout populations in the Malheur Recovery Unit.	69
APPENDIX F: List of Chapters	71

LIST OF TABLES

Table 1. Estimates of adult bull trout abundance in the North Fork Malheur, 1996 to 2001.	9
Table 2. List of bull trout streams on the 303d list that exceed the bull trout temperature standard.	12
Table 3. List of major forest fires in the upper Malheur River Basin since 1989	15
Table 4. Streams with riparian areas in poor condition in Upper Malheur River subbasin (USFS 2002).	17
Table 5. Chemical treatment projects in the Malheur Recovery Unit. Adapted from Bowers <i>et al.</i> (1993).	21

LIST OF FIGURES

Figure 1. Bull trout recovery units in the United States. The Malheur Recovery Units highlighted	2
Figure 2. Bull trout redd counts in the North Fork Malheur subbasin 1996 to present.	6
Figure 3. Map of the Malheur Recovery Unit with bull trout core area delineated.	31

INTRODUCTION

Recovery Unit Designation

Designation of the Malheur Recovery Unit is based in part on the inclusion of bull trout of this river basin within a single Gene Conservation Group by Oregon Department of Fish and Wildlife (Kostow 1995). The delineation of the Gene Conservation Groups is supported by the baseline genetic analysis conducted by Spruell and Allendorf (1997). Their analysis found that Malheur bull trout belong in the "Snake River" group of populations, but are distinct from other Oregon populations within this group. Further analysis by Spruell *et al.* (2002 in press) indicate Malheur bull trout are more genetically similar to bull trout populations from the Boise (Idaho) and Jarbidge (Nevada) drainages than to other populations in Oregon, and these three populations form a cluster within the Snake River group. Figure 1 shows the Malheur Recovery Unit.


Geographic Description

The Malheur River Basin is situated in eastern Oregon bordered on the south by the Owyhee River Basin, on the north by the Burnt River and John Day River Basins, on the west by the Malheur Lakes Basin, and by the Snake River to the east, which it enters near Ontario, Oregon. The basin includes portions of four counties; 62 percent occurs in Malheur County, 27 percent in Harney County, and small areas in Grant and Baker Counties (Malheur-Owyhee Watershed Council 1999).

From its headwaters in the Strawberry Range, at the southern terminus of the Blue Mountains, the Malheur River flows southeasterly for 105 kilometers (65 miles), turning north for 12 kilometers (19 miles), then east near the town of Juntura and continuing east to northeast to its confluence with the Snake River near the town of Ontario, a total distance of approximately 306 kilometers (190 miles) (Malheur-Owyhee Watershed Council 1999). Major tributaries include the South Fork Malheur River, which enters from the west at river kilometer 191 (river mile 119) near Riverside; the North Fork Malheur River, which enters at river kilometer 154 (river mile 96) near Juntura; Bully Creek, which enters at river kilometer 34 (river mile 21); and Willow Creek, which enters at river kilometer 32 (river mile 30). The latter three tributaries all enter the mainstem Malheur River from the north. The most important of the tributaries in terms of bull trout is the North Fork Malheur River, which also originates in the Strawberry Range and flows south to its confluence with the mainstem Malheur. Total drainage area of the Malheur River Basin is 11,940 square kilometers

(5,000 square miles) (Malheur-Owyhee Watershed Council 1999). Elevations in the basin range from the highest point on Graham

Figure 1. Bull trout recovery units in the United States. The Malheur Recovery Unit is highlighted.



Mountain at 2,613 meters (8,570 feet) to 610 meters (2,000 feet) at the mouth of the Malheur River (Hanson *et al.* 1990).

Public ownership accounts for approximately 66 percent of the land in the basin, most of it (47 percent) managed by the U.S. Bureau of Land Management, while 13 percent is managed by the Malheur National Forest and 6 percent is State-owned land. The remainder of the basin is in private or tribal ownership. Special management areas on the Malheur National Forest important to bull trout include the Strawberry Wilderness Area in the upper Malheur River, the

Monument Rock Wilderness Area in the North Fork Malheur River watershed, and the Wild and Scenic corridor in the North Fork Malheur River. In 2000, the Burns Paiute Tribe acquired 712 hectares (1,760 acres) in Logan Valley. The property encompasses approximately 11 kilometers (7 miles) of waterways, including portions of McCoy Creek, Big Creek, Lake Creek, Frazier Creek, and Malheur River (L. Schwabe, Burns Paiute Tribe, pers. comm. 2002).

Geology/Topography. Geologic processes that have shaped the Malheur River Basin include vulcanism, uplift, faulting, erosion, deposition, and to some extent glaciation. The Strawberry Range in the northwestern portion of the basin is composed primarily of Columbia River basalt dating from the Miocene but older outcrops from rocks dating to the Jurassic are also present (Franklin and Dyrness 1984). Subsequent uplifting of the basalt followed by faulting, erosion and weathering processes during the Pliocene and Pleistocene resulted in a varied relief of ridgetops, mountain slopes, dissected canyons and valley bottoms (Franklin and Dyrness 1984; U.S. Forest Service (USFS) 1999a). However, the bulk of the landscape is dominated by more gently sloping plateaus of composed of Miocene and Pliocene beds of tuffaceous sedimentary rocks capped by flows of rhyolite and basalt (Franklin and Dyrness 1984) and dissected by the stream network. At the lowest elevations near the mouth of the Malheur River, the basin is characterized by more gentle topography of low elevation terraces composed of lacustrine sediments and floodplains extending up the Malheur River and Willow Creek valleys (Malheur-Owyhee Watershed Council 1999).

Climate. The climate in the Malheur River Basin is continental, characterized by long, hot summers and cold winters. Summer high temperatures average between 85 and 95 degrees Celsius and winter temperatures average in the 20s. Annual precipitation averages from 25 to 30 centimeters (9.8 to 11.8 inches), with most occurring during winter as snow. Brief, intense storms occasionally occur during summer (Malheur-Owyhee Watershed Council 1999).

Hydrology. Flows in the Malheur River Basin are dominated by meltwater from the mountain snow pack. Peak discharge occurs in the spring (May through June) with traditional low flows in the summer and fall maintained by groundwater inflows. Springs originating in the Strawberry Range maintain year-round flows to streams they feed, while streams originating elsewhere have flows that tend to be ephemeral in nature (USFS 2000). Summer storms can influence streamflows with short duration, intense increases in runoff and streamflow.

The highly variable annual flows of the Malheur River and its tributaries have been harnessed through construction of storage and flood control facilities

and major diversion structures dating from 1881 with construction of the Nevada Diversion Dam at about river kilometer 31 (river mile 19) on the lower Malheur River. Winter flows are stored to provide reliable flows during the irrigation season, primarily to agricultural areas in the lower river valley. Major storage reservoirs include Beulah Reservoir (Agency Valley Dam constructed in 1934 at river kilometer 29 (river mile 18) on the North Fork Malheur River), Warm Springs Reservoir (Warm Springs Dam constructed in 1919 at river kilometer 198 (river mile 123) on the mainstem Malheur River, often referred to as the Middle Fork Malheur River above Warm Springs Reservoir) and Bully Creek (1962) and Malheur Reservoir (Willow Creek) in the lower Malheur basin. The entire flow of the Malheur River is diverted at Namorf Dam, river kilometer 111 (river mile 69), for re-distribution in the irrigation network of the lower Malheur basin. Irrigation return flows and groundwater inflows account for streamflow downstream of Namorf Dam. Namorf Dam (also known as Harper Dam) is laddered.

Vegetation. Vegetation in the Malheur Basin follows an elevation gradient from shrub-steppe to coniferous forest as the elevation and moisture increase. At the higher elevations the forest community includes lodgepole pine (*Pinus contorta*), whitebark pine (*P. albicaulis*), and subalpine fir (*Abies lasiocarpa*). Mixes of grand fir (*Abies grandis*), lodgepole pine, Douglas-fir (*Pseudotsuga menziesii*), western larch (*Larix occidentalis*), Engelmann spruce (*Picea engelmannii*) and ponderosa pine (*Pinus ponderosa*) occur on middle elevation slopes. The historical forest of ponderosa pine at the lower elevations now includes mixes of ponderosa pine, Douglas-fir, and grand fir due to the suppression of fire (USFS 1999b). Where the forest transitions to the grass and shrubland vegetation characteristic of the Great Basin province, communities of juniper (*Juniperus occidentalis*) and sagebrush (*Artemisia spp.*) dominate with mountain mahogany (*Cercocarpus spp.*) occupying rocky ridgetops (USFS 1999a). The native vegetation in the lower river valleys has been replaced by crops sustained through irrigation. The most common hardwood species found along riparian areas include alder (*Alnus spp.*), willow (*Salix spp.*), aspen (*Populus spp.*), and cottonwood (*P. trichocarpa*). However, long-term impacts from livestock grazing and fire suppression, and increased browsing pressure from expanding numbers of big game, have limited their occurrence and condition (USFS 1999b).

Cultural/Social. For thousands of years, the Wadatika band of the Northern Paiute Native Americans have inhabited the area that includes the Malheur River Basin and spans central and southeastern Oregon. Their seasonal migrations took them to the upper Malheur River to harvest salmon, as well as other native flora and fauna (Burns Paiute Tribe, *in litt.* no date; Malheur-Owyhee

Watershed Council 1999). Diseases introduced by early settlers greatly reduced the native population and altercations with nonnative settlers led to the establishment of the Malheur Reservation in 1873. The 719,758 hectares (1,778,560 acres) of the reservation included Castle Rock, Strawberry Mountain, the North and South Forks of the Malheur River, and portions of Harney County. Continued invasion of the Malheur Reservation by settlers led to the Bannock Indian War and eventual disbanding of the reservation in 1883. Descendants of the Wadatika band make up the current Burns-Paiute Tribe. They occupy a 312 hectare (771 acre) reservation near Burns, as well as property in Logan Valley.

Europeans first entered the basin in the 1820's when fur traders came in search of beaver (*Castor canadensis*). Settlers followed later in the century with the opening of the Oregon Trail and discovery of gold and silver in the Owyhee River Basin. Ontario is the largest urban area with a population of 10,000, followed by Vale with a population of 1,800 (Malheur-Owyhee Watershed Council 1999). Smaller unincorporated communities pertinent to the discussions of bull trout include Juntura and Drewsey.

Land Use. Agriculture is the dominant land use in the Malheur River Basin, with livestock production occurring over the greatest area. Irrigated agriculture dominates economics although it occupies only 4 percent of the landscape (Malheur-Owyhee Watershed Council 1999). Timber harvest and recreation occur in forested regions of the northwest part of the basin. The forest also provides summer livestock forage. Some mineral extraction occurs in the basin, primarily of diatomite, but deposits of gold and mercury cinnabar are also known to occur in the basin (Malheur-Owyhee Watershed Council 1999). Public ownership accounts for approximately 66 percent of land in the basin divided between the U.S. Forest Service (13 percent), Bureau of Land Management (47 percent), and State-owned land (6 percent) (Hanson *et al.* 1990).

Fish Species. Species found in association with bull trout in the North Fork Malheur River include redband/rainbow trout (*Onchorynchus mykiss*), bridgelip sucker (*Catostomus columbianus*), coarse scale sucker (*C. macrocheilus*), redband shiner (*Richardsonius balteatus*), mountain whitefish (*Prosopium williamsoni*), and northern pike minnow (*Ptychocheilus oregonensis*) (Schwabe *et al.* 2000). Species captured along with bull trout in the Middle Fork Malheur River during surveys in 2000, included brook trout (*Salvelinus fontinalis*), redband/rainbow trout, mountain whitefish, bridgelip suckers, speckled dace (*Rhinichthys osculus*), lognose dace (*R. cataractae*), sculpin (*Cottus spp.*), and redband shiner (*Richardsonius balteatus*) (Schwabe *et al.* 2001). A list of fish species found in the Malheur River Basin is presented in Appendix A.

DISTRIBUTION AND ABUNDANCE

Status of Bull Trout at the Time of Listing

Buchanan *et al.* (1997) classified bull trout in the North Fork Malheur River as "of special concern" and in the Upper Malheur River as at "high risk" of extinction. Categories of increasing extinction risk ranged from "low risk of extinction" to "probably extinct". Placement in each category was determined based on relative abundance, the severity of factors suppressing the population (for example, habitat conditions and presence of brook trout), and the potential of the population to recover to a healthy condition (Ratliff and Howell 1992). The category "of special concern" falls between a "low" and "moderate" risk level.

At the time of listing, the U.S. Fish and Wildlife Service considered all bull trout subpopulations in the Malheur Recovery Unit as "depressed" (63 FR 31647). Although subpopulations were an appropriate unit upon which to base the 1998 listing decision, the recovery plan has revised the biological terminology, to better reflect both current understanding of bull trout life history and conservation biology theory. Therefore, subpopulation terms will not be used in this chapter.

Current Distribution and Abundance

Bull trout are found in the North Fork Malheur River subbasin and in the upper Malheur River mainstem and tributaries upstream of the town of Drewsey. They are considered two distinct local populations because of their geographic isolation from construction of dams without fish passage on the mainstem Malheur River and North Fork Malheur Rivers.

North Fork Malheur River. Spawning and rearing takes place in the mainstem and tributaries upstream of Crane Crossing in the North Fork Malheur River. Results from studies of radio-tagged fish (18 tagged in Beulah Reservoir in 1999) showed that bull trout moved upstream from overwintering areas in Beulah Reservoir into the river from mid-April until late May in 1999 (Schwabe *et al.* 2000), and in 2000, some were observed in the river by mid-March (Schwabe *et al.* 2001). By June tagged fish were well distributed in the North Fork Malheur between Beulah Reservoir and the spawning areas. By early August the majority of tagged fish had moved upstream of Crane Creek confluence at river mile 42.8 (river kilometer 69) and some had moved into spawning tributaries by mid-July. The peak for migration into spawning tributaries occurred by mid to late-August. The peak in adult downstream

migration from spawning tributaries occurred in late September and bull trout returned to the reservoir between late October and mid-December (Schwabe *et al.* 2000 and 2001).

Spawning surveys were initiated in the North Fork Malheur subbasin in 1992 in streams with known or suspected bull trout populations (Buchanan *et al.* 1997). Based on data collected since 1992, bull trout spawning begins in late August and peaks in September. Redds have been observed as late as November (Schwabe *et al.* 2000). Spawning has been documented in the mainstem North Fork Malheur upstream of the mouth of Deadhorse Creek and in the following tributaries: Horseshoe Creek, Swamp Creek, Sheep Creek, Elk Creek, Crane Creek, and Little Crane Creek. Bull trout have been observed in Cow Creek during spawning surveys, but no redds have been found (Schwabe *et al.* 2000).

Subadult rearing and adult foraging occurs from the headwaters of the North Fork Malheur River down to, and in, Beulah Reservoir. In August of 1997, an interagency team of biologists snorkel surveyed the North Fork Malheur River from the confluence of the North Fork and Little Malheur Rivers upstream to the National Forest boundary. They documented bull trout rearing down to the confluence of the Little Malheur River. Sizes of bull trout ranged from 56 to 400 millimeters (2 to 16 inches) in length with the majority in the 100 to 200 millimeter (4 to 8 inch) size range. The largest bull trout observed was in the 300 to 400 millimeter (12 to 16 inch) size range (A. Miller, U.S. Forest Service, pers. comm. 2002). Trapping of subadult¹ bull trout during 1998 and 1999 using a rotary screw trap and passive integrated transponder (PIT) tags showed bull trout migrating downstream from spawning and rearing areas (upstream of Crane Creek) in the North Fork Malheur River. During the period the trap was operated (June to October) two peaks in migration were observed, the largest in June, and another smaller one in September. The smallest bull trout trapped in Beulah Reservoir during 1998 and 1999 measured 220 millimeters, or 8 to 9 inches in length (Schawbe *et al.* 2000).

Most radio-tagged bull trout overwintered in Beulah Reservoir. Some bull trout exit Beulah Reservoir during flood control operations, as well as during the irrigation season and are lost to the population above the dam. The extent of use and survival of bull trout in the mainstem Malheur River downstream of Agency Dam is unknown. During the 1999 study, five bull trout


¹

Fish were classified as subadult based on scale analysis that showed most fish to be age 3 or 4. Age 3 fish ranged in fork length from 168 to 260 millimeters (7 to 10 inches), while age 4 fish ranged in fork length from 214 to 315 millimeters (8 to 12 inches) (Schwabe *et al.* 2000).

were fitted with radio tags and released below the dam. Most of the radio-tagged bull trout stayed within 1.5 kilometers (1 mile) of the dam during the study period (Schwabe *et al.* 2000).

Redd surveys are used to track bull trout trends in abundance in the Malheur River Basin. A general upward trend in redd numbers has been observed since 1992. Redd counts from 1996 to the present are shown in Figure 2. Survey areas and timing have been standardized since 1996, so these data represent the most accurate estimate of trend available (Tinneswood and Perkins 2001).

Figure 2. Bull trout redd counts in the North Fork Malheur subbasin 1996 to present.



In 2000, surveyors observed 29 redds with 2 or more bull trout on them, or 19 percent of the total redds counted (151). The number of bull trout observed on a single redd ranged from one to five, or approximately 2.4 bull trout per redd (Tinneswood and Perkins 2001). This represents the maximum bull trout per redd ratio recorded for the North Fork Malheur River local population. In 2001, when 29 redds and bull trout were counted, the ratio was closer to 2.0 (R. Perkins, Oregon Department of Fish and Wildlife, pers. comm. 2002). These ratios along with redd counts for the period 1996 to 2001 were used to estimate

abundance of adult bull trout in the North Fork Malheur (Table 1). An estimate of 4,132 bull trout age at least one year old for the North Fork Malheur River is based on population sampling completed in 1991 and 1992 using a multiple pass removal method (Buchanan *et al.* 1997).

Table 1. Estimates of adult bull trout abundance in the North Fork Malheur, 1996 to 2001.

Year	Redds	Range of Abundance
1996	38	76 to 91
1997	64	128 to 154
1998	74	148 to 178
1999	115	230 to 276
2000	153	306 to 367
2001	125	250 to 300

Upper Malheur River. Bull trout spawning and juvenile rearing occurs in the Upper Malheur River and tributaries upstream of the confluence of Big Creek. Streams where redds have been identified include Snowshoe Creek, Meadow Fork Big Creek, Lake Creek, Summit Creek, and Big Creek, although brook trout may account for some of the redds.

Timing of bull trout spawning in the Upper Malheur population is similar to what has been observed in the North Fork Malheur population with the peak occurring in mid-September. Data collected in 1999 showed that 40 percent of the redds were counted prior to September 15th. These redds were assumed to be bull trout redds as they occurred in streams where most of the bull trout were also observed, although brook trout were present during surveys (Schwabe *et al.* 2000).

Subadult rearing and adult foraging occurs downstream to approximately river kilometer 286 (river mile 178) in the vicinity of Hog Flat, based on limited historical and recent radio-telemetry documentation (Schwabe *et al.* 2001), and one radio-tagged fish was tracked to near the mouth of Wolf Creek in the spring of 2002 (W. Bowers, Oregon Department of Fish and Wildlife, pers. comm. 2002). It is possible, although not documented, that fish forage as far downstream as Warm Springs Reservoir during winter.

A general upward trend in redds counted in the Upper Malheur River has been observed for the period of record, 1998 to present. However, an estimate of adult abundance for the Upper Malheur River local population is not available at this time because of the inability to distinguish between bull trout and brook trout redds when not occupied.

Oregon Department of Fish and Wildlife estimated the bull trout population at 3,554 bull trout at least one year old, based on population sampling in 1993 and 1994 in Big Creek, Lake Creek, and the Meadow Fork of Big Creek. Densities ranged from a high of 0.474 fish per lineal meter (762.8 fish per mile) in Meadow Fork of Big Creek to a low of 0.039 fish per lineal meter (62.8 fish per mile) in Lake Creek (Buchanan *et al.* 1997).

REASONS FOR BULL TROUT DECLINE

Threats to bull trout arise from the modification or destruction of their habitat or from direct take (e.g., harvest, disease, injury, etc.). Fish habitat in the Malheur Recovery Unit has been altered significantly since European settlement, affecting not only bull trout, but anadromous species as well. Salmon and steelhead are considered an important part of the historical prey base for bull trout. Historical land uses affecting bull trout habitat in the Malheur Basin include livestock grazing, timber harvest, road building, dispersed recreation, and irrigated agriculture. Liberal harvest regulations and fish stocking programs have also been implicated in the decline of bull trout (Buchanan *et al.* 1997).

Prior to the construction of dams, bull trout in the Malheur River had access to the Snake River, although their typical summer habitat was in the upper part of the basin. The lower reaches of the Malheur River are considered too warm in the summer for bull trout rearing and spawning, but did serve as a migration corridor to the Snake and Columbia Rivers, and as foraging habitat during winter. At the time of listing, the U.S. Fish and Wildlife Service identified forestry, grazing, agricultural practices, water quality, and introduced species (brook trout) as threats to Malheur River bull trout (63 FR 31647).

Subwatersheds in the Malheur River Basin were assessed by the Malheur-Owyhee Watershed Council using the Proper Functioning Condition methodology developed by the Bureau of Land Management (1995) to evaluate the streams based on their capability and potential to collect, store, and release water in a beneficial manner. Based on subbasin assessments, the mainstem Malheur River upstream of Warm Springs Dam was judged to be in fair condition with some channel incision and bank erosion upstream of Warm Springs Reservoir, primarily in the Drewsey and Logan Valleys. In these areas the channel has been altered to facilitate agriculture and to some extent road building (Malheur-Owyhee Watershed Council 1999). Only the upper reaches in the North Fork Malheur River Subbasin have been assessed thus far, and they were judged to be in fair to good condition. Evidence of streambank erosion in some areas between Beulah Reservoir and Crane Creek may be natural or resulting from activities occurring upstream (Malheur-Owyhee Watershed Council 1999). Downstream of Beulah Dam the North Fork Malheur River has been channelized in one section and altered by agriculture and road building in two other sections (Malheur-Owyhee Watershed Council 1999). The Malheur River from Warm Springs Dam downstream to the mouth is, "incised along much of its length with exposed actively eroding streambanks" (Malheur-Owyhee Watershed Council 1999) and constrained for much of its length by a highway and railroad bed.

Temperature is the most serious water quality parameter affecting bull trout in the Malheur Basin based on the Department of Environmental Quality 303d list (Table 2). Sediment loading has been observed in the North Fork Malheur River both upstream and downstream of Beulah Reservoir and downstream of Warm Springs Dam during spring runoff and local storm events (Malheur-Owyhee Watershed Council 1999).

Table 2. List of bull trout streams on the 303(d) list that exceed the bull trout temperature standard.

(Source: <http://www.deq.state.or.us/wq/WQLData/SubBasinList98.asp>)

Waterbody Name	Boundaries
Big Creek	Mouth to Meadow Fork
Crane Creek	Mouth to Little Crane Creek
Elk Creek	Mouth to Headwaters
Lake Creek	Mouth to Headwaters
Little Crane Creek	Mouth to Headwaters
Malheur River, North Fork	Crane Creek to Headwaters
Sheep Creek	Mouth to Headwaters
Swamp Creek	Mouth to Headwaters

Other parameters listed for stream segments downstream of currently occupied bull trout habitat include bacteria, chlorophyll-a, and toxics (<http://www.deq.state.or.us/wq/WQLData/SubBasinList98.asp>). Most of the water quality issues have resulted from both historical and current land use activities in the basin. However, summer stream temperatures were likely limiting due to natural conditions, especially during hot, dry periods with low precipitation.

Dams

Construction and operation of Agency Dam/Beulah Reservoir and Warm Springs Dam and reservoir has fragmented Malheur Basin bull trout local populations, altered the stream temperature and streamflow regimes, halted migration of anadromous species and their nutrient inputs, altered forage bases, and in the case of Agency Dam, entrained bull trout during spring and summer (U.S. Bureau of Reclamation (USBOR) 1998, Hanson *et al.* 1990; Schwabe *et al.*

2000). Releases of water in the spring are through the spillway radial gates. Four of 39 bulltrout radio-tagged above Beulah Reservoir were entrained through Agency Dam (Schwabe *et al.* 2000).

Neither dam was constructed with fish passage, nor were conservation pools included for fish. During drought years there is insufficient water to hold overwintering bull trout in either reservoir, although use by bull trout has not been documented in Warm Springs Reservoir. The reduction in reservoir volume also affects production of aquatic organisms and forage species important to bull trout, although the prey base would be more concentrated and more vulnerable to predation. At low reservoir levels, bull trout are likely entrained through the outlet works and are lost to the population (USBOR 1998). The extent of entrainment through the dam outlet works and the resulting effect on the adfluvial bull trout population is unknown at this time.

At the end of the irrigation season releases from both dams cease. Fish that survive entrainment are stranded in pools downstream. Data from a few fish radio-tagged in the pool below the dam show they remained within 2 kilometers (about 1 mile) of the location where they were tagged (Schwabe *et al.* 2000). During the winter dams do not release water so the reservoirs can refill. Lack of flows in the North Fork Malheur downstream of Beulah Reservoir is an issue from October 15 until spring when the reservoir fills and spills or irrigation season begins. In the absence of entrainment data for Agency Dam, the U. S. Fish and Wildlife Service estimates the expected level of take to be at least seven percent of the total population present in the reservoir. This is based on the entrainment rate determined at Arrowrock Dam (on the Boise River in Idaho), which was the only entrainment data available for a Bureau of Reclamation facility (USFWS 1999).

Habitat below dams is deprived of gravel and wood inputs that are trapped behind the dams. Streambank stability and riparian habitat may be negatively affected by extended surges in flows released during peak irrigation demands (Malheur-Owyhee Watershed Council 1999).

Agency Valley Dam and Beulah Reservoir have created an environment which supports adfluvial bull trout. These fish are probably larger than they would have been historically because of the abundant food supply present in the reservoir environment. Water temperatures downstream of Beulah are probably cooler than they were historically because the releases come from the deepest portion of the reservoir where water is coolest (R. Rieber, *in litt.* 2000).

Forest Management Practices

The following discussion pertains primarily to public lands as there is little data available on forest management practices on private land. Best Management Practices recommended by the Oregon Department of Forestry under the Oregon Forest Practices Act (ORS 527.610 to 527.770, 527.990(1) and 527.992) and Forest Practice Administrative Rules (OAR 629.600 to 629.665) guide forest management on private lands. Forest management practices on the Malheur National Forest affecting bull trout habitat include harvest of forest products, associated road building, and recreation activities. Livestock grazing and hay production also take place in the forested zone on public and private land, and these will be discussed in subsequent sections.

A proper functioning condition analysis for bull trout watersheds on the National Forest in the North Fork Malheur subbasin determined that streams were "functioning at unacceptable risk" and did not meet criteria for temperature, sediment, substrate embeddedness, pool frequency and quality, and large pools (USFS 1999b). A similar analysis for bull trout watersheds in the upper Malheur subbasin determined streams were "functioning at unacceptable risk" and did not meet criteria for temperature, sediment, physical barriers, pool frequency and quality, and road density and location (USFS 1999a).

The natural disturbance regime in the bull trout occupied subbasins includes some flooding associated with rain-on-snow events and wildfires ignited by late summer dry thunderstorms. Wildfires present the greatest natural threat due to conditions resulting from historical fire exclusion, management of stands in dense multi-story character, and accumulations of fuels associated with insect and disease related mortality (USFS 1999a; USFS 1999b). The Glacier, Sheep Mountain, and Snowshoe fires may have directly affected bull trout streams by increasing temperature and sedimentation (Buckman *et al.* 1992). Stream temperature data from Swamp Creek showed average maximum temperatures above 10 degrees C at sites within the fire boundary with very little vegetative cover and below 10 degrees C at the site outside the fire boundary (Perkins 2000). Major forest fires and acreage burned in recent history are shown in Table 3. Although some monitoring of sediments has occurred, additional monitoring of sediment in North Fork Malheur subbasin is needed to assess the impact of the Glacier, Sheep Mountain, and Snowshoe fires on bull trout habitat.

North Fork Malheur River. The mainstem of the North Fork Malheur River on the Malheur National Forest lacks instream wood and large wood for recruitment, as well as side channel habitat and channel complexity. Above Crane Crossing this is attributed to past fires, commercial logging, and personal

use wood cutting. Lack of large wood contributes to lack of pools, and loss of channel complexity, bank stability, and shade. Reduced sinuosity in unconstrained forested reaches outside the wilderness is attributed to lack of large wood associated with historical harvest activity (USFS 1999b).

Table 3. List of major forest fires in the upper Malheur River Basin since 1989 (USFS 1999a and USFS 1999b). Area burned is in acres/hectares

Subbasin	Fire	Year	Area Burned
Upper Malheur	Sheep Mountain	1990	2,076 / 840
Upper Malheur	Snowshoe	1990	9,285 / 3,757
North Fork Malheur	Glacier	1989	3,237 / 1,310
North Fork Malheur	Ironside	1994	5,261 / 2,129
North Fork Malheur	Monument Rock	1989	2,098 / 849
North Fork Malheur	Powder	1994	5,806 / 2,350
North Fork Malheur	Sheep Mountain	1990	8,867 / 3,588

In addition to bull trout spawning and rearing segments that exceed the bull trout temperature standard (see previous Water Quality section), temperature is also limiting in the North Fork Malheur from Crane Creek to Beulah for summer subadult and adult rearing due in part to forest management practices and irrigation withdrawals. This segment is on the 303(d) list for not meeting Oregon water quality standard for cold water species (64 degrees F) (<http://www.deq.state.or.us/wq/WQLData/SubBasinList98.asp>).

The road system used for timber harvest on National Forest lands within the Malheur Recovery Unit dates from the mid-1950's. Until that time a series of railroad spur lines were used to haul logs to mills in Burns and Seneca (USFS 1999b). Roads constructed along stream channels have narrowed the potential for stream channel movement (decreased sinuosity) and reduced vegetation potential in riparian areas. Twenty-six roads (16 miles) occur within Riparian Habitat Conservation Areas, a management designation whereby riparian areas are protected and enhanced for wildlife, non-anadromous fish habitat, and water quality values (USFS 1999b).

Bull trout occupied streams surveyed in the upper North Fork Malheur River show fine sediment estimates ranging from 31 percent to 40 percent (Oregon Department of Fish and Wildlife data cited in USFS 1999b). Additional studies in bull trout spawning and rearing areas should be done to isolate

sediment sources, which may include roads and burned areas (e.g., Glacier Fire), and to define and prioritize remedial action.

Upper Malheur River. Passage barriers are known to exist in the upper Malheur River subbasin at road crossings on Summit Creek and at the old railroad grade crossing on Bosonberg Creek (USFS 2000). A number of road culverts on the National Forest present passage barriers as well. A forest-wide culvert inventory was conducted in the summer of 2001. Preliminary analysis showed that many culverts would not pass juveniles under low flow conditions, although the data has not been sorted for barriers to bull trout (A. Miller, pers. comm. 2002).

Seasonal thermal barriers at the mouths of Lake, McCoy and Summit Creeks limit bull trout movement between local populations and access to potential habitat (USFS 2002). These are areas where stream gradients decrease as the streams enter natural open meadow areas. However, water diverted from streams on the National Forest to irrigate hay meadows on private land has decreased the amount of water in the streams, increasing the potential for warming (A. Miller, pers. comm. 2002).

Inputs of fine sediments from surface erosion have been attributed to timber harvest, grazing (both livestock and big game), and wildfire (USFS 1999a). Sediment in Snowshoe and Corral Creek Basins may be related to fire salvage activities following the Snowshoe Fire in 1990, while high levels of fine sediment in Lake Creek and Bosonberg Creek are road-related (USFS 2002). Contributing to the road-related sedimentation on the National Forest may be the high road densities (2.4 to 2.6 miles/square mile) where road densities are greater than stream densities (USFS 2000). This is indicative of the potential for road surfaces to alter the way water reaches the stream by increasing the runoff rate and decreasing infiltration rates. Thirty-one kilometers (19 miles) of roads are in Riparian Habitat Conservation Areas (USFS 1999a). Trail crossings on the Sheep Creek and Crane Creek trails are experiencing increased degradation of habitat (USFWS 2001), presumably from sediment loading among other impacts.

The lack of sufficient large pools may be limiting production of bull trout in the Upper Malheur subbasin. Large wood in the stream channel plays an important role in pool formation, and although large wood is considered adequate overall (USFS 2002), Lake Creek and Summit Creek wood levels are below Inland Native Fish Strategy standards (USFS 1995, USFS 1999a). Lack of pools has also been attributed to the extirpation of beaver by the mid-1850's and subsequent loss of pools created by beaver dams (USFS 2000).

Areas that need riparian restoration are identified in Table 4. Bluebucket Creek and associated tributaries has also been identified as needing riparian and stream restoration (USFS 1999a). Areas that burned in the Snowshoe and Corral Basin fires are still recovering and lack adequate shade (USFS 2002).

Livestock Grazing

Livestock grazing is a major land use activity on public and private lands in the North Fork Malheur River and Upper Malheur River subbasins. Livestock are turned

out onto public grazing allotments in the spring and summer, while private pastures are used for hay production for winter forage. Livestock are gathered in the fall and winter in riparian pastures, which in addition to trampling riparian vegetation, may also contribute to poor water quality. Impacts to bull trout habitat resulting from over-utilization of the riparian vegetation include loss of shade, which can increase stream temperature; increased bank erosion through removal of woody species and physical trampling of streambanks; and direct trampling of redds where livestock have access to spawning areas.

While much of the impact is from historical grazing practices and corrective measures have been taken, some problem areas remain on public and private lands. Unauthorized use is a continuing problem in both U.S. Forest Service and Bureau of Land Management allotments (USFWS 2001; U.S. Bureau of Land Management (BLM) 2002). Recently, wild ungulates (deer and elk) have also been implicated in the suppression of riparian vegetation (D. Young, *in litt.* 2000; USFWS 2001; BLM 2002).

Table 4. Streams with riparian areas in poor condition in Upper Malheur River subbasin (USFS 2002).

Stream Name	Ownership	Reach
Big Creek	Private	Mouth to Forest Service Road 16
Big Creek	U.S. Forest Service	Small reaches from Forest Service Rd 1648 to Wilderness Boundary
Malheur	Private, U.S. Forest Service	Malheur Ford to Big/Lake confluence
Snowshoe Creek	U.S. Forest Service	Upper reach (Snowshoe Fire)

Table 4. Streams with riparian areas in poor condition in Upper Malheur River subbasin (USFS 2002).

Stream Name	Ownership	Reach
Corral Basin Creek	U.S. Forest Service	Mouth to headwaters
Bosonberg Creek	Private	Mouth to Forest Service Rd 16
Bosonberg Creek	U.S. Forest Service	Upper reach (Snowshoe Fire)
Lake Creek	Private, U.S. Forest Service	Mouth to Forest Service Rd 16
McCoy Creek	Private	Mouth to FS Boundary
Summit Creek	U.S. Forest Service, Private	Mouth to North end of Summit Prairie
Crooked Creek	U.S. Forest Service	Mouth to Headwaters

North Fork Malheur River. Lack of riparian habitat from grazing is a limiting factor in the North Fork Malheur River downstream of Crane Crossing to Beulah Reservoir and from Agency Dam to the mouth of the river, primarily on the private lands. Vegetation is routinely removed by ranchers to facilitate livestock operations (Malheur-Owyhee Watershed Council 1999). During spawning surveys on Elk Creek from 1997 to 2000, cattle were present (Tinnewood and Perkins 2001). Presence of sheep in Swamp Creek and cattle in Little Crane Creek, both important bull trout spawning tributaries, was noted during 1999 spawning surveys (Schwabe *et al.* 2000).

Upper Malheur River. Lack of riparian habitat is a problem in the Malheur River between Malheur Ford and Bosonberg Creek and in the lower reaches of Bosonberg, Big, Lake, Corral Basin, and McCoy Creeks. Streams on private lands in Logan Valley and Summit Prairie have altered stream channels and flows as a result of livestock grazing and water withdrawals (USFS 1999a). Riparian and stream channel restoration is needed on Bluebucket Creek and its tributaries (USFS 1999a). Sedimentation from grazing is a problem in the Malheur River downstream to the Drewsey Valley. Downstream of Highway 20 the channel is aggrading. This could be due to excessive sediment inputs upstream of the highway from bank erosion as a result of poor grazing practices.

Agricultural Practices

Water is diverted from both forks of the Malheur River for stock water and irrigation of hay meadows. Many of the diversions originating on the National Forest are held by private landowners and are small spring sources for domestic water sources and irrigation (USFS 1999a). Not all diversions have been inventoried. From the mouth of the North Fork Malheur River to the confluence with the Snake River, the Malheur River flows through intensively managed agricultural lands where water quality and water quantity issues magnify. The Malheur River downstream of Namorf Dam to the mouth is not currently considered suitable migration habitat for bull trout because of water quality and passage issues.

Stream temperature may be affected by diversions as well as by surface releases from Beulah Reservoir. Diverted water reduces available instream habitat can result in increased stream temperature, and also creates migration barriers.

Where diversions are unscreened, fish can become entrained in the irrigation ditches and perish. There are four unscreened diversions in the North Fork Malheur River subbasin upstream of Beulah Reservoir and six or more downstream of the dam. Some of the diversions above the Drewsey Valley in the upper Malheur River subbasin are screened, however none are screened from the upper Drewsey Valley downstream. Some irrigation diversions in the Drewsey Valley are also passage barriers. The diversion at Namorf Dam is not screened, nor are any of the diversions downstream to the mouth of river screened (R. Perkins, pers. comm. 2002). During the early spring virtually all of the river is diverted at Namorf Dam into the canal to fill Bully Creek Reservoir (W. Bowers, pers. comm. 2002).

Transportation Network

Most of the threats associated with the transportation network occur on the National Forest and were discussed previously under Forest Management Practices. On tribal land within the Malheur National Forest the access ford to Burn Paiute Tribe property in Logan valley is unstable and contributes excess sediment to Lake Creek. A bridge originally accessed the property, but it collapsed in 1999. The Burns Paiute Tribe's goal for restoration is to stabilize the stream banks and reduce sediments to Lake Creek by installing culverts designed to handle a 50-year flood event.

Highway 20 follows the river corridor through the canyon downstream of Juntura. Should the area be used by bull trout in the future there would always be a threat from spills of toxic materials transported on the highway.

Mining

Mining was not identified as an issue historically or currently.

Residential Development

Residential development was not identified as an issue historically or currently.

Fisheries Management

Historical fishery management with its liberal bag limits on trout (5 to 10 trout per day bag limit depending on location) were a factor in the bull trout's decline in the Malheur Basin. Currently, poaching and incidental mortality associated with sport fisheries may be a limiting factor in the recovery unit. Bull trout that are entrained at Agency Dam and remain in the plunge pool are vulnerable to harvest.

Another practice that may have impacted bull trout was the chemical treatment to eliminate fish from many stream segments upstream of Beulah and Warm Springs Reservoirs. Some bull trout mortalities were observed (Bowers *et al.* 1993). These areas were then planted with hatchery rainbow trout. Table 5 lists chemical treatment projects in the basin.

The fish toxicant most often used is rotenone, a natural substance derived from the roots of several South American plants. It acts by entering the blood stream of the fish through the gills and preventing oxygen use at the cellular level. In addition to fish, benthic invertebrates, zooplankton, and to a lesser extent, amphibians, are susceptible to rotenone. Mammals, birds, and plants are not directly affected, but may be influenced indirectly by the removal of fish and other organisms from the biological community. Mammals and birds can drink treated water without adverse effects (California Department of Fish and Game 1985).

Introduced brook trout are a limiting factor in the upper Malheur River where bull trout x brook trout hybrids have been identified (the North Fork Malheur does not contain brook trout). Brook trout compete with bull trout for food, cover, and spawning areas and hybridize with bull trout. Gunckel (2000)

Table 5. Chemical treatment projects in the Malheur Recovery Unit. Adapted from Bowers *et al.* (1993).

Subbasin	Year	Areas treated
North Fork Malheur	1950	Beulah Reservoir
	1955	North Fork Malheur River tributaries and Beulah Reservoir
	1961	Beulah Reservoir and tributaries, mainstem Malheur River from mouth of Crane Creek to Beulah Reservoir
	1968	Beulah Reservoir and lower sections of tributaries, Little Malheur River
	1977	North Fork Malheur River 8 kilometers above Beulah Reservoir and the reservoir
	1987	North Fork Malheur River from Castle Rock Ranch downstream to mouth near Juntura, Beulah Reservoir
Upper Malheur	1955	Malheur River from Dollar Basin ford downstream; Lake Creek, McCoy Creek; upper Crooked Creek; Big Creek, Bosonberg Creek; Summit Creek; other tributaries to Warm Springs Reservoir and selected reservoirs including Warm Springs

researched feeding behavior and diet of bull trout and brook trout and found that interference competition was likely due to their similar habitat use, feeding behavior and diet, and aggressive interactions between the species. The dominant behavior of brook trout place bull trout at a disadvantage when resources are scarce (Gunckel 2000). In addition, rainbow trout have been stocked in the past in the vicinity of campgrounds on public land. These fish also would have competed with bull trout for food, shelter, and space.

Unauthorized introductions of nonnative fish and other aquatic organisms are always a concern and are difficult to prevent. Bait fish are often accidentally introduced, while some introductions of game fish are intentional. The most recent unauthorized introduction occurred in Beulah Reservoir when crappie (*Pomoxis spp.*) were discovered in 2001. Impacts to the bull trout population are

unknown at this time. Efforts will be undertaken by Oregon Department of Fish and Wildlife to remove them to prevent their establishment in the reservoir.

There are no significant fish disease issues in the recovery unit at this time. Disease was not identified as threat to bull trout when they were listed. Whirling disease has been present since the 1980's in the Grande Ronde Basin, which also has several local populations of bull trout. However, bull trout there do not exhibit signs of the disease. Periodic sampling for whirling disease is a necessary fish management tool to monitor and detect any spread of the disease.

Bull trout may be inherently resistant to some diseases that are more devastating to other salmonids. In challenge studies conducted by Oregon State University researchers, Metolius (Deschutes) bull trout exposed to high and low doses of the infectious stages of *Myxobolus cerebralis* (causative agent in whirling disease) showed no signs of infection as measured by presence of spores, clinical disease signs, or histopathology. Rainbow trout exposed simultaneously showed high infection prevalence and disease severity. Nor were infections detected in Metolius (Deschutes) bull trout exposed to infection by *Ceratomyxosis shasta* (Bartholomew 2001). Disease studies conducted on bull trout from the Deschutes River Basin showed them to be relatively resistant to all strains of Infectious Hematopoietic Necrosis Virus tested. Bull trout had detectable levels of antigen to *R. salmoninarum* (bacterial kidney disease) but no evidence of the disease.

Isolation and Habitat Fragmentation

As previously discussed in the "Dams" section, the two populations in the Malheur River Basin are isolated from one another by the Agency and Warm Springs Dams. Other passage barriers exist that prevent gene flow among local populations within the two subbasins and inhibit expansion of local populations into potential bull trout habitat (*i.e.*, culverts, diversions, and thermal barriers). Losses to the local populations when bull trout are entrained at Agency Dam and in unscreened irrigation diversions or through natural catastrophic events, such as drought and wildfire, have impacted bull trout numbers, although losses are difficult to quantify. When a bull trout population size is small such losses increase the probability of extinction (Rieman and McIntyre 1993). Recovery and long-term persistence of bull trout in the Malheur Recovery Unit is dependent on reconnecting the two isolated local populations. Therefore, the most critical challenges facing Malheur Basin bull trout are eliminating the threat of brook trout in the upper Malheur subbasin and providing opportunities for passage at Agency and Warm Springs Dams.

ONGOING RECOVERY UNIT CONSERVATION MEASURES

Efforts to recover bull trout and fish habitat in general are ongoing in the Malheur Recovery Unit with a high level of cooperation between fishery entities on various projects. For example, the Oregon Department of Fish and Wildlife, the Burns Paiute Tribes, U.S. Bureau of Land Management, U.S. Forest Service, and the Bureau of Reclamation staff work have worked cooperatively on bull trout spawning and habitat surveys for many years. The Malheur River Basin has several active local watershed groups and conservation districts dedicated to finding workable solutions to restoring watershed health. The following list is by no means complete, but is representative of ongoing efforts within the recovery unit.

Oregon Department of Fish and Wildlife

The Oregon Department of Fish and Wildlife has reduced or eliminated hatchery rainbow and brook trout stocking programs; adopted changes in angling regulations to prohibit take of bull trout in 1992, modified regulations on other fisheries to reduce incidental take; made changes to in-water work periods to better address bull trout needs; and has applied for Instream Water Rights on ten stream segments to benefit bull trout (Appendices C and D). Stream priorities for additional acquisition of instream water rights are also shown in Appendices C and D.

The agency has also developed and distributed bull trout identification posters to provide better public awareness education. Large metal signs were installed in 2000 at major entry points in the recovery unit where people might encounter bull trout.

A multi-year bull trout research project was initiated by the Oregon Department of Fish and Wildlife in northeastern Oregon in 1995 to study bull trout life history, ecology, and genetics. Funding has been provided through a grant from the Bonneville Power Administration. Part of the research project examined brook trout and bull trout interactions in the Middle Fork Malheur. Oregon Department of Fish and Wildlife also has obtained funding through a cooperative agreement with the U.S. Fish and Wildlife Service (section 6) funding to assist with spawning surveys to monitor bull trout populations.

Planning efforts in the Oregon Department of Fish and Wildlife have focused on formation of local bull trout working groups to develop conservation strategies for Oregon bull trout populations. This effort was begun in 1993 and these working groups were later used to form the foundation for bull trout

recovery unit teams in Oregon. Bull trout distribution mapping using the agency's Geographic Information System began in 1996. The statewide bull trout assessment, Status of Oregon's Bull Trout (Buchanan *et al.* 1997), was published in 1997.

Oregon State Police

Bull trout remain a high priority for enforcement through the Oregon State Police Cooperative Enforcement Program. Staff from Oregon State Police and Oregon Department of Fish and Wildlife meet annually to set priorities for enforcement in the Malheur Recovery Unit through the Cooperative Enforcement Program.

Bureau of Reclamation

The Bureau of Reclamation is actively engaged in a number of studies resulting from a U.S. Fish and Wildlife Service biological opinion for operation and maintenance of Bureau of Reclamation projects (*i.e.*, Beulah and Warm Springs Reservoirs) in the recovery unit. Investigations are focused on water quality, safety of dams, water acquisitions, conservation pools, and fish passage. The Bureau of Reclamation has initiated or committed to the following studies.

In August 1997, the Bureau of Reclamation entered into a cooperative agreement with the U.S. Forest Service, Bureau of Land Management, Oregon Department of Fish and Wildlife, and the Burns Paiute Tribe to investigate the distribution and abundance of bull trout in the North Fork Malheur River basin from Beulah Reservoir to the headwaters. The study involves radio telemetry for up- and downstream migrating juvenile and adult bull trout and a continuation of surveys of bull trout spawning grounds. In 1999, bull trout were captured below Agency Valley Dam, radio-tagged, and monitored to determine the potential for entrainment through the dam.

Water quality monitoring/modeling studies were initiated in 1999 by the Bureau of Reclamation to determine the minimum pool elevation necessary to support adfluvial bull trout. Bi-monthly field sampling for water quality and limnology study was conducted in 1999 and 2000. The data, along with a bioenergetics model being developed for bull trout by the U.S. Geological Survey, will be used to develop a reservoir model to quantitatively define the effects of various pool elevations on dissolved oxygen depletion and winter kill during the winter months, and to determine the available habitat for bull trout during the summer thermal stratification period when existing data suggest

oxygen depletion in the deeper waters and high temperatures at the surface (R. Rieber, U.S. Bureau of Reclamation (BOR), pers. comm. 2002).

In 2000, the Bureau of Reclamation started releasing up to 650 cubic feet per second (18 cubic meters per second) of water through the outlet works from the bottom of Beulah reservoir and not the spillway to test if this will reduce entrainment of bull trout. The spillway is only operated if releases greater than 650 cubic feet per second (18 cubic meters per second) are required as a result of flood control operations, and those are generally of short duration.

The Bureau of Reclamation has investigated alternatives for creating a minimum fisheries pool. The Beulah Reservoir Conservation Pool Appraisal study looked at three options: raising capacity at Beulah, raising capacity at Warm Springs, or building a new reservoir at Vines Canyon (near Vale) and exchanging water in it for a conservation pool at Beulah. Storing additional water behind Warm Springs Dam would be the least expensive of the options evaluated in this report if this work can be coordinated with the modification planned for Warm Springs Dam (USBOR 2001).

An appraisal-level investigation on possible upstream passage or trapping options at Agency Valley Dam is scheduled to begin in 2003, pending results from the water quality research (R. Rieber, pers. comm. 2002).

U.S. Fish and Wildlife Service

The U.S. Fish and Wildlife Service is engaged in ongoing section 7 consultations with the U.S. Forest Service, the Bureau of Land Management, and the Bureau of Reclamation. Biological opinions issued by the U.S. Fish and Wildlife Service cover grazing, vegetation management, and reservoir operations of the respective agencies.

The U.S. Fish and Wildlife Service has provided funding through the Partners for Fish and Wildlife Program to install fish screens above Drewsey. They have also partnered with the Natural Resources Conservation Service, the National Fish and Wildlife Foundation, Oregon Department of Fish and Wildlife, Oregon Watershed Enhancement Board, and private landowners to implement a variety of projects that will directly benefit bull trout habitat. These projects include fish screens, modification of water diversions, riparian fencing, water conservation projects, and other actions to improve instream and passage conditions. This area has been identified as a geographic priority area under the Environmental Quality Incentive Program under the U.S. Department of Agriculture Farm Bill.

U.S. Forest Service and U.S. Bureau of Land Management

The U.S. Forest Service and Bureau of Land Management have taken steps to improving grazing management in the recovery unit. Changes in season of use, setting utilization standards, fencing some areas to prevent livestock access to the streams, and increased monitoring of use and enforcement action when necessary are all strategies used to improve riparian habitat and thus improve conditions for bull trout.

The U.S. Forest Service closed their diversion on Lake Creek and is considering turning it into an instream water right. Other measures to protect and improve stream habitat over the past 20 years include road closures, riparian exclosures, hardwood planting and caging, slash riprap along streambanks, and log weir placement (USFS 1999a and 1999b).

RELATIONSHIP TO OTHER CONSERVATION EFFORTS

State of Oregon

On January 14, 1999, Governor Kitzhaber expanded the Oregon Plan for Salmon and Watersheds (State of Oregon 1997) to include all at-risk wild salmonids throughout the State through Executive Order 99-01. The goal of the Oregon Plan is to, "restore populations and fisheries to productive and sustainable levels that will provide substantial environmental, cultural, and economic benefits." Components of this plan include (1) coordination of efforts by all parties, (2) development of action plans with relevance and ownership at the local level, (3) monitoring progress, and (4) making appropriate corrective changes in the future. It is a cooperative effort of State, local, Federal, tribal and private organizations, and individuals.

Oregon Department of Fish and Wildlife and Oregon Water Resources Department have established priorities for restoration of streamflow as part of the Oregon Plan for Salmon and Watersheds (Measure IV.A.8). The Oregon Department of Fish and Wildlife has prioritized streamflow restoration needs by ranking biophysical factors, water use patterns, and the extent that water limits fish production in a particular area. Oregon Water Resources Department watermasters will incorporate the priorities into their field work activities as a means to implement flow restoration measures. The needs priorities will be used by the Oregon Watershed Enhancement Board as one criterion in determining funding priorities for enhancement and restoration projects. Watershed councils and other entities may also use the needs priorities as one piece of information to determine high priority restoration projects. Bull trout occupied streams in the recovery unit are included in the highest priority designation for streamflow restoration (NWPPC 2002).

Opportunities to convert existing out-of-stream flows to instream flows in Oregon are available through a variety of legislatively mandated programs administered by Oregon Water Resources Department, such as transfers of type and place of use (ORS 536.050(4)), voluntary written agreement among water users to rotate their use of the supply to which they are collectively entitled (ORS 540.150 and OAR 690-250-0080), allocation of "conserved water" to instream use (ORS 537.455 to 537.500), lease of all or a portion of consumptive water rights to instream purposes (ORS 537.348, OAR 690-77-070 to 690-77-077), exchange of a water right for an instream purpose to use water from a different source, such as stored water, surface, or ground water (ORS 540.533 to 540.543), and substitute a ground water right for a primary surface water right (ORS

540.524). Oregon Water Trust provides purchase of water rights from willing landowners for conversion to instream water rights.

Through the Malheur River Basin Total Maximum Daily Load process a water quality management plan will be developed to address forest, agricultural, urban, and transportation sources of water quality impairment. Monitoring is scheduled to begin in the summer of 2003, which will likely delay development of the Total Maximum Daily Load scheduled for that same year (D. Butcher, Oregon Department of Environmental Quality, pers. comm. 2002). (See also <http://waterquality.deq.state.or.us/wq/TMDLs/TMDLs.htm> for more information.)

The Agricultural Water Quality Management Program, established through the Senate Bill 1010 process (ORS 568.900 through 568.933), addresses water pollution associated with agricultural lands and activities. The Malheur River Basin Agricultural Water Quality Management Area Plan was drafted in 2001. Major areas of concern include (1) pollution control and waste management (excessive concentrations of nutrients, particularly nitrogen and phosphorus, in surface water), (2) sediments carried in irrigation return flows, (3) riparian area management to restore vegetation, (4) improved streambank stability, and (5) rangeland and pasture management to improve water infiltration rates. Actions to address resource concerns will be through voluntary efforts of individual landowners with technical assistance provided to landowners through the Natural Resources Conservation Service (NRCS). A monitoring plan will be implemented to characterize baseline conditions, track plan implementation and evaluate plan effectiveness (ODA 2001).

Burns Paiute Tribe

The Burns Paiute Tribe has obtained funding through the Bonneville Power Administration Fish and Wildlife Program for a multi-year research project to learn more about the life history of bull trout and other native salmonids in the Malheur River Basin. The project is part of a cooperative effort toward bull trout recovery that includes participation by other members of the Malheur Recovery Unit Team. The Tribe has also acquired several ranch properties (with Bonneville Power Administration mitigation funding and assistance from The Nature Conservancy) that include sections of the North Fork, South Fork and Malheur River mainstem. The ranch properties will be restored to improve habitat for fish and wildlife (Northwest Power Planning Council 2002).

Local Planning Efforts

There are two active watershed councils in the Malheur River Basin, the Malheur-Owyhee Watershed Council and the Bully Creek Watershed Council. The Malheur-Owyhee Watershed Council was formed in 1995, with the mission "To lead the effort to conserve, protect and enhance all watershed resources for optimum economic and environmental benefits within the Malheur watershed". They completed the Malheur Basin Action Plan in 1999 to provide local guidance and solutions for meeting State and Federal mandates, primarily the Clean Water Act and the Endangered Species Act (Malheur-Owyhee Watershed Council 1999).

Both the Harney and Malheur Soil and Water Conservation Districts work with local private landowners in partnership with the Natural Resources and Conservation Service and Farm Services Agency to implement conservation activities on private land.

Northwest Power Planning Council's Subbasin Planning

As part of the Pacific Northwest Electric Power Planning and Conservation Act of 1980, the Bonneville Power Administration has the responsibility to protect, mitigate and enhance fish and wildlife resources affected by operation of Federal hydroelectric projects in the Columbia River and tributaries. The Northwest Power Planning Council develops and coordinates the Columbia River Basin Fish and Wildlife Program that is implemented by the Bonneville Power Administration, U.S. Army Corps of Engineers, U.S. Bureau of Reclamation, and Federal Energy Regulatory Commission. Coordination of Bonneville Power Administration's responsibilities for protection, enhancement, and mitigation and incorporation of recommendations by the Northwest Power Planning Council is in part accomplished through the development of subbasin summaries, which identify the status of fish and wildlife resources, limiting factors, and recommended actions at the subbasin level.

The Draft Malheur Subbasin Summary (NWPPC 2002) encompasses the Malheur Recovery Unit, and is consistent with bull trout recovery planning efforts to identify limiting factors. The draft subbasin summary identifies dams, unscreened irrigation diversions, reduced instream flows, poor water quality, loss of shade and channel structure and function, nonnative species, sediment from roads, past fisheries management (including liberal harvest limits, stocking of nonnative trout, and chemical treatment projects), loss of beaver and beaver dam complexes, and extirpation of salmon as contributing to the decline of bull trout.

The overall fisheries goal of the Draft Malheur Subbasin Plan is, "to protect, enhance and restore where needed, resident and anadromous fish in their historical habitat". The Malheur Recovery Unit Team will continue to utilize this planning process to identify and seek funding for projects to aid bull trout recovery.

STRATEGY FOR RECOVERY

A core area represents the closest approximation of a biologically functioning unit. The combination of core habitat (*i.e.*, habitat that could supply all the necessary elements for the long-term security of bull trout including both spawning and rearing as well as foraging, migrating, and overwintering) and a core population (*i.e.*, bull trout inhabiting a core habitat) constitutes the basic core area upon which to gauge recovery within a recovery unit. Within a core area, many local populations may exist.

Malheur Core Area. For purposes of recovery, the Malheur Recovery Unit contains one core area, the Malheur Core Area, which encompasses tributaries containing two local populations (and additional potential local populations as identified by the recovery unit team) and the mainstem Malheur River from headwaters downstream to Namorf Dam. Although bull trout would have had seasonal access to the Malheur River downstream of Namorf Dam, there is no spawning

Figure 3. Map of the Malheur Recovery Unit with bull trout core area delineated.



and juvenile rearing habitat downstream of this point. Migration and overwinter suitableing/foraging habitat downstream to Namorf Dam is sufficient to support recovered populations.

The Malheur Core Area (Figure 3) includes two local populations located in the headwaters of the North Fork Malheur River and the Upper Malheur River subbasins. Additional research on homing fidelity may indicate further division into additional local populations is appropriate.

Recovery Goals and Objectives

The goal of the bull trout recovery plan is to ensure the long-term persistence of self-sustaining, complex, interacting groups of bull trout distributed throughout the species' native range, so the species can be delisted. To achieve this goal the following objectives have been identified for bull trout in the Malheur Recovery Unit:

Maintain the current distribution of bull trout within the core area and re-establish bull trout in previously occupied habitats in the Upper Malheur River and tributaries and the North Fork Malheur River and tributaries.

- Maintain stable or increasing trends in abundance of bull trout in the Malheur Recovery Unit. This will require increasing abundance within the two local populations (Upper Malheur and North Fork Malheur).
- Restore and maintain suitable habitat conditions for all bull trout life history stages and strategies.
- Conserve genetically diverse local populations of bull trout within the Malheur Recovery Unit by providing opportunities for genetic exchange between the local populations. This can best be achieved by ensuring connectivity between the North Fork Malheur River and the Upper Malheur River.

Recovery criteria for the Malheur Recovery Unit reflect the stated objectives and consideration of population and habitat characteristics within the recovery unit. The Malheur Recovery Unit Team evaluated the current status of bull trout based on four population elements. The four elements were: (1) number of local populations, (2) adult abundance (defined as the number of sexually mature fish present in a core area in a given year), (3) productivity (defined as a measure of population trend and variability), and (4) life history forms (as an indicator of the functional connectivity of the system). For each element, the Malheur Recovery Unit Team classified bull trout based on relative risk categories.

Rieman and McIntyre (1993) and Rieman and Allendorf (2001) evaluated the bull trout population numbers and habitat thresholds necessary for long-term viability of the species. They identified four elements, and the characteristics of those elements, to consider when evaluating the viability of bull trout populations. These four elements are (1) number of local populations; (2) adult abundance (defined as the number of spawning fish present in a core area in a given year); (3) productivity, or the reproductive rate of the population (as measured by population trend and variability); and (4) connectivity (as represented by the migratory life history form and functional habitat). For each element, the Malheur Recovery Unit Team classified bull trout into relative risk categories based on the best available data and the professional judgment of the team.

The Malheur Recovery Unit Team also evaluated each element under a potential recovered condition to produce recovery criteria. Evaluation of these elements under a recovered condition assumed that actions identified within this chapter had been implemented. Recovery criteria for the Malheur Recovery Unit reflect (1) the stated objectives for the recovery unit, (2) evaluation of each population element in both current and recovered conditions, and (3) consideration of current and recovered habitat characteristics within the recovery unit. Recovery criteria will probably be revised in the future as more detailed information on bull trout population dynamics becomes available. Given the limited information on bull trout, both the level of adult abundance and the number of local populations needed to lessen the risk of extinction should be viewed as a best estimate.

This approach to developing recovery criteria acknowledges that the status of populations in some core areas may remain short of ideals described by conservation biology theory. Some core areas may be limited by natural attributes or by patch size and may always remain at a relatively high risk of extinction. Because of limited data within the Malheur Recovery Unit, the recovery unit team relied heavily on the professional judgment of its members.

Local Populations. Metapopulation theory is important to consider in bull trout recovery. A metapopulation is an interacting network of local populations with varying frequencies of migration and gene flow among them (Meffe and Carroll 1994) (Chapter 1). Multiple local populations distributed and interconnected throughout a watershed provide a mechanism for spreading risk from stochastic events. In part, distribution of local populations in such a manner is an indicator of a functioning core area. Based in part on guidance from Rieman and McIntyre (1993), bull trout core areas with fewer than 5 local populations are at increased risk, core areas with between 5 and 10 local populations are at intermediate risk, and core areas with more than 10 interconnected local populations are at diminished risk.

For the Malheur Core Area, there are currently 2 known local populations (Upper Malheur River and North Fork Malheur River). Based on the aforementioned guidance, the Malheur Core Area is at increased risk from stochastic events.

Adult Abundance. The recovered abundance levels in the Malheur Recovery Unit were determined by considering theoretical estimates of effective population size, historical census information, and the professional judgment of recovery team members. In general, effective population size is a theoretical concept that allows us to predict potential future losses of genetic variation within a population due to small

population sizes and genetic drift (Chapter 1). For the purpose of recovery planning, effective population size is the number of adult bull trout that successfully spawn annually. Based on standardized theoretical equations (Crow and Kimura 1970), guidelines have been established for maintaining minimum effective population sizes for conservation purposes. Effective population sizes of greater than 50 adults are necessary to prevent inbreeding depression and a potential decrease in viability or reproductive fitness of a population (Franklin 1980). To minimize the loss of genetic variation due to genetic drift and to maintain constant genetic variance within a population, an effective population size of at least 500 is recommended (Franklin 1980; Soule 1980; Lande 1988). Effective population sizes required to maintain long-term genetic variation that can serve as a reservoir for future adaptations in response to natural selection and changing environmental conditions are discussed in Chapter 1 of the recovery plan.

For bull trout, Rieman and Allendorf (2001) estimated that a minimum census number of 50 to 100 spawners per year was needed to minimize potential inbreeding effects within local populations. Furthermore, a census population size between 500 and 1000 adults in a core area is needed to minimize the deleterious effects of genetic variation due to drift.

For the purposes of bull trout recovery planning, abundance levels were conservatively evaluated at the local population and core area levels. Local populations containing fewer than 100 spawning adults per year were classified as at risk from inbreeding depression. Bull trout core areas containing fewer than 1,000 spawning adults per year were classified as at risk from genetic drift

From 1996 to 2001, annual spawner survey estimates in the North Fork Malheur local population have averaged approximately 95 redds per year. Based on this information, this local population is not considered at risk from inbreeding depression. Limited data on the Upper Malheur River local population precluded

evaluation of inbreeding risks. Overall, the Malheur Core Area most likely contains less than 1,000 spawning adults and is considered at risk from the deleterious effects of genetic drift.

Productivity. A stable or increasing population is a key criterion for recovery under the requirements of the Endangered Species Act. Measures of the trend of a population (the tendency to increase, decrease, or remain stable) include population growth rate or productivity. Estimates of population growth rate (*i.e.*, productivity over the entire life cycle) that indicate a population is consistently failing to replace itself, indicate increased extinction risk. Therefore, the reproductive rate should indicate the population is replacing itself, or growing.

Since estimates of the total population size are rarely available, the productivity or population growth rate is usually estimated from temporal trends in indices of abundance at a particular life stage. For example, redd counts are often used as an index of a spawning adult population. The direction and magnitude of a trend in the index can be used as a surrogate for the growth rate of the entire population. For instance, a downward trend in an abundance indicator may signal the need for increased protection, regardless of the actual size of the population. A population which is below recovered abundance levels but moving toward recovery would be expected to exhibit an increasing trend in the indicator.

The population growth rate is an indicator of probability of extinction. This probability cannot be measured directly, but it can be estimated as the consequence of the population growth rate and the variability in that rate. For a population to be considered viable, its natural productivity should be sufficient for the population to replace itself from generation to generation. Evaluations of population status will also have to take into account uncertainty in estimates of population growth rate or productivity. For a population to contribute to recovery, its growth rate must indicate that the population is stable or increasing for a period of time.

Connectivity. The presence of the migratory life history form within the Malheur Recovery Unit was used as an indicator of the functional connectivity of the recovery unit and both core areas. If the migratory life form was absent, or if the migratory form is present but local populations lack connectivity, the core area was considered to be at increased risk. If the migratory life form persists in at least some local populations, with partial ability to connect with other local populations, the core area was judged to be at intermediate risk. Finally, if the migratory life form was present in all or nearly all local populations, and had the ability to connect with other local populations, the core area was considered to be at diminished risk.

Migratory bull trout persist in the Malheur River and at least partial connection exists between local populations within the core area and were considered at intermediate risk.

Recovery Criteria

Recovery criteria for bull trout in the Malheur Recovery Unit are the following:

1. **Bull trout are distributed among 2 or more local populations in the recovery unit within the Malheur Core Area.** In a recovered condition one or more local populations would occur within the Upper Malheur River and one or more local populations would occur within the North Fork Malheur River. There is potential to further separate the Upper Malheur River into 2 or 3 local populations, and the North Fork Malheur River into 2 or more local populations. However, additional population studies and a better understanding

of bull trout fidelity to their natal streams is needed to further define local populations in the recovery unit. There may be potential to expand the current distribution of bull trout into additional tributaries within their historic range, such as Little Malheur River (North Fork Malheur tributary) and Crooked Creek, Bosonberg Creek, McCoy, and Corral Basin (Upper Malheur River tributaries). The identified recovered distribution may place the Malheur Core Area at increased risk from stochastic events. Natural habitat features within the Malheur River Basin may limit expansion of bull trout distribution to idealized levels identified in this chapter. After additional population and genetic information is collected, recovery criteria may be revised, and the risk level associated with stochastic events re-evaluated.

2. **Estimated abundance of adult bull trout in the Malheur Recovery Unit is between 2,000 and 3,000 individuals distributed between the two local populations.** The recovery unit team expects to achieve this criteria by securing the current population and increasing the abundance of migratory adults in Upper Malheur River. The recovered abundance range was derived using the professional judgement of the recovery unit team and estimation of productive capacity of identified local populations. This abundance level would mean that the core area and local populations would not be at risk from inbreeding or genetic drift, respectively. These goals may be refined as more information becomes available, through monitoring and research.
3. **Adult bull trout exhibit a stable or increasing trend for at least 2 generations at or above the recovered abundance level within the**

Malheur Core Area. The development of a standardized monitoring and evaluation program which would accurately describe trends in bull trout abundance is identified as a priority research need. As part of the overall recovery effort, the U.S. Fish and Wildlife Service will take the lead in addressing this research need by forming a multi-agency technical team to develop protocols to evaluate trends in bull trout populations.

4. **Specific barriers inhibiting bull trout movement and recovery in the Malheur Recovery Unit have been addressed, ensuring opportunities for connectivity among local populations within the core area.** This means addressing passage at Agency Dam on the North Fork Malheur, Warm Springs Dam on the Malheur River, all unscreened diversions in core areas, and all impassable culverts. Additional studies will be needed to determine the feasibility of providing two-way passage at Beulah and Warm Springs Dams. Reduction or elimination of threats from brook trout interaction in the Upper Malheur will need to be accomplished prior to restoration of passage to ensure the success of bull trout recovery. While the major connectivity issues in the

Malheur Recovery Unit are associated with Agency Dam and Warm Springs Dam, additional gains in recovery of bull trout populations through expansion of habitat within the two subbasins can be achieved by restoring passage over and around barriers at road crossings, culverts, and water diversions. The known barriers are listed in Appendix B and specific action to address them are highlighted in the recovery narrative portion of this plan. There may be others that have not been identified.

Recovery criteria for the Malheur Recovery Unit were established to assess whether recovery actions have resulted in the recovery of bull trout. The Malheur Recovery Unit Team expects that the recovery process will be dynamic and require refinements as more information becomes available over time. While removal of bull trout as a species under the Endangered Species Act (*i.e.*, delisting) can only occur for the entity that was listed (Columbia River Distinct Population Segment), the criteria listed above will be used to determine when the Malheur Recovery Unit Recovery Unit is fully contributing to recovery of the population segment.

Research Needs

Based on the best scientific information available, the recovery unit team has identified recovery criteria and actions necessary for recovery of bull trout within the Malheur Recovery Unit. However, the recovery unit team recognizes

that many uncertainties exist regarding bull trout population abundance, distribution, and recovery actions needed. The recovery unit team feels that if effective management and recovery are to occur, the recovery plan for the Malheur Recovery Unit should be viewed as a "living" document, to be updated as new information becomes available. As part of this adaptive management approach, the Malheur Recovery Unit Team has identified essential research needs within the recovery unit.

Monitoring and Assessment. The Malheur Recovery Unit Team based estimates of recovered abundance levels and number of local populations on the best available information and professional judgement. Information on historical abundance levels and distribution of spawning populations is limited. The recovery unit team realizes that recovery criteria will most likely be revised as recovery actions are implemented and bull trout populations begin to respond. The recovery unit team will rely on adaptive management to better refine both abundance and distribution criteria. Adaptive management is a continuing process of planning, monitoring, evaluating management actions, and research. This adaptive management approach will identify actions that maximize the ability to achieve recovery objectives. In addition, this approach will provide a better understanding of key uncertainties, crucial to long-term management actions.

Monitoring and evaluation of population levels and distribution are important components of any adaptive management approach. The U.S. Fish and Wildlife Service will take the lead in developing a comprehensive monitoring approach that will provide guidance and consistency in evaluating bull trout populations. An important component in recovery implementation and the use of adaptive management will be the evaluation of recommended actions.

The Malheur Recovery Unit Team has identified an urgent need for the development of a standardized monitoring and assessment program that would more accurately describe current status of bull trout within the recovery unit, as well as identify improvements in current sampling protocols that would allow for monitoring the effectiveness of recovery actions. Development and application of models that assess population trend and extinction risk will be useful in refining recovery criteria as the recovery process proceeds.

Additional research needed for recovery include studies that evaluate effectiveness of restoration techniques, limiting factors analysis in watersheds with historical bull trout habitat and potential habitat, identification of sediment sources and impacts to bull trout, and continuation of life history studies.

Genetic Studies. The Malheur Recovery Unit Team recommends that studies be initiated to describe the genetic makeup of bull trout in the core area. Genetic information on local populations within the core area is necessary for a more complete understanding of bull trout interactions and population dynamics, and may lead to revision of recovery criteria. In addition, a recovery unit-wide evaluation of the current and potential threat of bull trout hybridization with brook trout is needed. The ability to evaluate the potential harm to specific local populations could be used in prioritizing management actions. Genetic baseline information would also be a necessity in the implementation of any artificial propagation program.

ACTIONS NEEDED

Recovery Measures Narrative

In this chapter and all other chapters of the bull trout recovery plan, the recovery measures narrative consists of a hierarchical listing of actions that follows a standard template. The first-tier entries are identical in all chapters and represent general recovery tasks under which specific (*e.g.*, third-tier) tasks appear when appropriate. Second-tier entries also represent general recovery tasks under which specific tasks appear. Second-tier tasks that do not include specific third-tier actions are usually programmatic activities that are applicable across the species' range; they appear in *italic type*. These tasks may or may not have third-tier tasks associated with them; see Chapter 1 for more explanation. Some second-tier tasks may not be sufficiently developed to apply to the recovery unit at this time; they appear in *a shaded italic type (as seen here)*. These tasks are included to preserve consistency in numbering tasks among recovery unit chapters and intended to assist in generating information during the comment period for the draft recovery plan, a period when additional tasks may be developed. Third-tier entries are tasks specific to the Malheur Recovery Unit. They appear in the implementation schedule that follows this section and are identified by three numerals separated by periods.

The Malheur Recovery Unit chapter should be updated or revised when recovery tasks are accomplished, environmental conditions change, or monitoring results or other new information becomes available. Revisions to the Malheur Recovery Unit chapter will likely focus on priority streams or stream segments within core areas where restoration activities occurred, and habitat or bull trout populations have shown a positive response. The Malheur Recovery Unit Team should meet annually to review annual monitoring reports and summaries, and make recommendations to the U.S. Fish and Wildlife Service.

- 1 Protect, restore, and maintain suitable habitat conditions for bull trout.
 - 1.1 Maintain or improve water quality in bull trout core areas or potential core habitat.
 - 1.1.1 Assess sediment sources in Malheur Basin affecting bull trout. Identify road-related sediment problem areas in the Malheur Core Area. Examine the ways roads capture and channel runoff, and changes in surface runoff associated with soil compaction. Identify the source of stream channel

aggradation in the Malheur River downstream of Highway 20.

- 1.1.2 Stabilize roads, crossings, and other sources of sediment delivery. Address sediment issues identified in the upper Malheur River watershed assessment, (e.g., road related sediment in Lake and Bosonberg Creeks) and as a result of task 1.1.1. Correct sedimentation at the ford that accesses the Burns Paiute Tribe property in Logan Valley. Construction of a culvert crossing (several culverts) that will sustain a 50 year flood event has been proposed.
- 1.1.3 Assess and mitigate effects on bull trout from nonpoint source pollution. Temperature and sedimentation are the most pressing nonpoint source pollution issues affecting bull trout. Data collection will be used for designation of Total Maximum Daily Loads.
- 1.1.4 Increase monitoring of sediment inputs on the Malheur National Forest. Additional monitoring is needed to assess sediment to bull trout spawning and rearing habitat resulting from wildfires in the Malheur Core Area (Snowshoe and Corral Basins).
- 1.2 Identify barriers or sites of entrainment for bull trout and implement tasks to provide passage and eliminate entrainment.
 - 1.2.1 Identify and implement opportunities for two-way passage at major dams. Agency and Warm Springs Dams and the Drewsey diversion are priorities.
 - 1.2.2 Provide passage at transportation/road-related barriers identified in completed and ongoing surveys. Oregon Department of Transportation surveys have been completed for State and County roads in Oregon. The U.S. Forest Service completed a culvert inventory on the Forest in 2001, but analysis has not been completed as barriers pertain to bull trout. Establish priorities for passage provision and implement necessary actions.

- 1.2.3 Install appropriate fish screens and passage structures around diversions and/or remove related migration barriers. High priorities for screening include diversion on Big Creek that conveys water to the Burns Paiute Tribal property (Oxbow Ranch) on the upper Malheur River and the Castle Rock diversions on the North Fork Malheur.
- 1.2.4 Restore connectivity and opportunities for migration by improving instream flows. Use the Oregon Department of Fish and Wildlife instream flow priorities as a guide. See list in Appendices B, C, D, and E.
- 1.3 Identify impaired stream channel and riparian areas and implement tasks to restore their appropriate functions.
 - 1.3.1 Restore shade and canopy, riparian cover, and native vegetation in all bull trout spawning areas. Summit Creek, the Malheur River downstream of Logan Valley, and Crooked Creek in the Upper Malheur subbasin have suppressed woody vegetation in reaches that need this component to become stable, as well as tributary streams to the North Fork Malheur on the lower end of the Malheur National Forest.
 - 1.3.2 Reduce grazing impacts in all bull trout spawning areas. Implement measures to reduce livestock impacts to streams, (e.g., fencing, changes in timing and use of riparian pastures, off-site watering and salting), to accomplish this task. Areas that would benefit from shade restoration include lower Lake Creek, lower Big Creek, lower Bosonberg Creek, lower McCoy Creek, Crane Creek, and Buttermilk Creek, as well as, private and tribal lands in Logan Valley. The highest grazing management priority for the U.S. Fish and Wildlife Service is that no direct effects occur to spawning fish or redds after August 15.
 - 1.3.3 Review habitat information to identify and prioritize opportunities for channel restoration in Lake Creek. Design and implement projects based on findings. Lake Creek is an important bull trout spawning stream that is impacted by

sediment from roads, the ford on Tribal land, and is lacking large wood.

- 1.3.4 Improve instream habitat in the Malheur Core Area. Implement INFISH guidelines (USFS 1995). Use existing habitat surveys or new surveys if necessary to identify opportunities to improve habitat complexity and restore channel morphology.
- 1.4 Operate dams to minimize negative effects on bull trout in reservoirs and downstream.
 - 1.4.1 Review reservoir operational concerns and provide operating recommendations. Improve where needed, water level manipulation, methods of release, entrainment, minimum fisheries pool, and fish passage. Reservoirs of highest concern are Beulah and Warm Springs Reservoirs.
 - 1.4.2 Establish and provide instream flows downstream from reservoirs and stabilize flow regimes. Improve flows in the winter downstream of Beulah Reservoir from October 15 until the irrigation season begins in the spring, to improve the suitability of this reach as overwintering habitat for bull trout that pass through or over Agency Dam.
- 1.5 Identify upland conditions negatively affecting bull trout habitats and implement tasks to restore appropriate functions.
 - 1.5.1 Assess current and historical effects of upland management on changes to the hydrograph. (e.g., timing and magnitude of peak flows) in all spawning tributaries. Restore vegetation in forested portions of the Malheur Core Area in areas where bull trout spawning and juvenile rearing occur.
- 2 Prevent and reduce negative effects of nonnative fishes and other nonnative taxa on bull trout.
 - 2.1 *Develop, implement, and enforce public and private fish stocking policies to reduce stocking of nonnative fishes that affect bull trout.*

- 2.2 *Evaluate enforcement policies for preventing illegal transport and introduction of nonnative fishes.*
- 2.3 *Increase education and outreach to the public about ecosystem concerns of illegal introductions of nonnative fishes.*
 - 2.3.1 Develop and implement an educational effort to address problems and consequences of unauthorized fish introductions. Curtail illegal introductions of crappie in Beulah Reservoir (discovered in 2001).
- 2.4 *Evaluate biological, economic, and social effects of control of nonnative fishes.*
- 2.5 *Implement control of nonnative fishes where found to be feasible and appropriate.*
 - 2.5.1 Implement brook trout removal effort(s) wherever feasible and biologically supportable. Prioritize streams in the upper Malheur River that contain brook trout x bull trout hybrids for removal effort and implement action.
- 2.6 *Develop tasks to reduce negative effects of nonnative taxa on bull trout.*
- 3 *Establish fisheries management goals and objectives compatible with bull trout recovery, and implement practices to achieve goals.*
 - 3.1 *Develop and implement State and tribal native fish management plans integrating adaptive research.*
 - 3.1.1 Incorporate bull trout recovery actions into The Oregon Plan for Salmon and Watersheds and the Pacific Northwest Power Planning Council Subbasin plans. Request assistance with implementation of recovery strategies for bull trout through both planning processes.
 - 3.1.2 Coordinate bull trout recovery with recovery efforts, management plans, etc. of other species, such as redband trout, in the Malheur Core Area.

- 3.2 Evaluate and prevent over harvest and incidental angling mortality of bull trout.
 - 3.2.1 Maintain bull trout as high priority for Oregon's Cooperative Enforcement Program.
 - 3.2.2 Increase information outreach to anglers. Provide information on bull trout identification, special regulations, methods to reduce hooking mortality of bull trout caught incidentally, and the value of bull trout and their habitat and their place in the ecosystem.
 - 3.2.3 Improve and implement fisheries management guidelines and policies designed to protect native species. Some examples include the Oregon Draft Native Fish Conservation Policy and the Malheur River Basin Fish Management Plan.
- 3.3 Evaluate potential effects of introduced fishes and associated sport fisheries on bull trout recovery and implement tasks to minimize negative effects on bull trout.
 - 3.3.1 Determine site-specific levels of competition and hybridization with introduced sport fish and assess impacts of those interactions. Assess severity of threat due to hybridization with brook trout in the Upper Malheur local population.
- 3.4 *Evaluate effects of existing and proposed sport fishing regulations on bull trout.*
- 4 Characterize, conserve, and monitor genetic diversity and gene flow among local populations of bull trout.
 - 4.1 Incorporate conservation of genetic and phenotypic attributes of bull trout into recovery and management plans.
 - 4.1.1 Develop a genetic management plan. Develop a genetic management plan for the Malheur Recovery Unit including the establishment of genetic baselines for each local population, monitoring genetic changes in existing local

populations, determination of new local populations, and identification of actions needed to maintain existing opportunities for gene flow among bull trout populations.

- 4.2 *Maintain existing opportunities for gene flow among bull trout populations.*
- 4.3 *Develop genetic management plans and guidelines for appropriate use of transplantation and artificial propagation.*
- 5 Conduct research and monitoring to implement and evaluate bull trout recovery activities, consistent with an adaptive management approach using feedback from implemented, site-specific recovery tasks.
 - 5.1 Design and implement a standardized monitoring program to assess the effectiveness of recovery efforts affecting bull trout and their habitats.
 - 5.1.1 Evaluate effectiveness of different habitat restoration techniques used in restoring stream functions and local bull trout populations in the Malheur Core Area.
 - 5.2 Conduct research evaluating relationships among bull trout distribution and abundance, bull trout habitat, and recovery tasks.
 - 5.2.1 Identify site-specific threats that may be limiting bull trout in watersheds with historical bull trout habitat. Use the list in Appendix E to prioritize work.
 - 5.2.2 Determine the movement and seasonality of use of different habitat types by adult and sub-adult migratory bull trout in multiple streams, with emphasis on reservoirs (e.g., Beulah and Warm Springs) and mainstem rivers in the Malheur Core Area.
 - 5.2.3 Investigate potential for restoring historic prey base by reintroducing anadromous species, and take action based on findings. The Burns Paiute Tribe has a proposal before the Northwest Power Planning Council to assess the feasibility of the upper Malheur watershed to support reintroduction of

anadromous population above Beulah and Warm Springs reservoirs (NWPPC 2002).

- 5.3 *Conduct evaluations of the adequacy and effectiveness of current and past best management practices in maintaining or achieving habitat conditions conducive to bull trout recovery.*
- 5.4 *Evaluate effects of diseases and parasites on bull trout, and develop and implement strategies to minimize negative effects.*
- 5.5 Develop and conduct research and monitoring studies to improve information concerning the distribution and status of bull trout.
 - 5.5.1 Review and update databases for bull trout distribution records. Including StreamNet (2002), which provides data and data services for fish in the Pacific Northwest.
 - 5.5.2 Conduct surveys in potential habitat in the Malheur Core Area where bull trout status is unknown or recolonization is anticipated.
 - 5.5.3 Determine life history requirements of resident and migratory bull trout local populations in the Malheur Core Area. Knowledge of specific requirements of bull trout in this recovery unit will facilitate their management.
 - 5.5.4 Determine consequences of genetic fragmentation and isolation due to human-made barriers. Examples include Agency and Warm Springs Dams, and impassable diversions and culverts.
 - 5.5.5 Evaluate food web interactions in drainages most affected by introduced fishes, reservoir operations, loss of anadromous species (prey base/nutrients), etc. Studies are underway in Beulah Reservoir to study the predator/prey relationship between bull trout and other species in the reservoir. Some research has been completed on brook trout bull trout interaction in the Upper Malheur subbasin.

- 5.6 *Identify evaluations needed to improve understanding of relationships among genetic characteristics, phenotypic traits, and local populations of bull trout.*
- 6 Use all available conservation programs and regulations to protect and conserve bull trout and bull trout habitats.
 - 6.1 Use partnerships and collaborative processes to protect, maintain, and restore functioning core areas for bull trout.
 - 6.1.1 Provide long-term habitat protection. Explore opportunities to protect bull trout habitat through land purchase from willing sellers, conservation easements, management plans, land exchanges, etc. Promote collaborative efforts to establish or support existing local watershed groups and Soil and Water Conservation Districts to accomplish site specific protection/restoration activities. Priority reaches in the Malheur Recovery Unit that would benefit from this action include the North Fork Malheur between Beulah Reservoir and the National Forest boundary, Bosonberg Creek, Lake Creek, McCoy Creek, Big Creek, and Crane Creek.
 - 6.1.2 Work cooperatively with the Burns-Paiute tribal government to implement recovery actions.
 - 6.1.3 Develop educational materials on bull trout and their habitat needs to provide to landowners and interested public parties. Some examples include information on watershed form and function, riparian and channel restoration, and large wood placement.
 - 6.1.4 Identify and pursue opportunities to implement recovery strategies. Garner support from management agencies, pursue cooperative funding, partnerships, challenge cost share opportunities, and other private and governmental grants; and utilize mitigation and natural resource damage settlement funds as available.
 - 6.1.5 Integrate watershed analyses and assessments and restoration activities on public and private lands. Current restoration initiatives include the Total Maximum Daily Load process

and Senate Bill 1010 water quality management plans, watershed council activities, and Federal watershed analyses. Sites that would benefit from this action include Summit and Crooked Creeks in the Upper Malheur River.

- 6.2 *Use existing Federal authorities to conserve and restore bull trout.*
- 6.3 *Enforce existing Federal, State, and Tribal habitat protection standards and regulations and evaluate their effectiveness for bull trout conservation.*
- 7 Assess the implementation of bull trout recovery by recovery units, and revise recovery unit plans based on evaluations.
 - 7.1 Convene annual meetings of each recovery unit team to review progress on recovery plan implementation.
 - 7.1.1 Develop an annual work plan to support implementation in the Malheur Recovery Unit.
 - 7.2 *Assess effectiveness of recovery efforts.*
 - 7.3 Revise scope of recovery as suggested by new information.
 - 7.3.1 Periodically review progress towards recovery goals and assess recovery task priorities. Annually review progress toward population and adult abundance criteria and recommend changes, as needed, to the Malheur Recovery Unit chapter. In addition, review tasks, task priorities, completed tasks, budget, time frames, particular successes, and feasibility within the Malheur Recovery Unit.

IMPLEMENTATION SCHEDULE

The Implementation Schedule that follows describes recovery task priorities, task numbers, task descriptions, duration of tasks, potential or participating responsible parties, total cost estimate and estimates for the next five years, if available, and comments. These tasks, when accomplished, are expected to lead to recovery of bull trout in the Malheur Recovery Unit. Cost estimates are not provided for tasks which are normal agency responsibility under existing authorities. It should be noted that many of the estimated costs assigned to these recovery tasks will also benefit listed salmon and steelhead.

Parties with authority, responsibility, or expressed interest to implement a specific recovery task are identified in the Implementation Schedule. Listing a responsible party does not imply that prior approval has been given or require that party to participate or expend any funds. However, willing participants will benefit by demonstrating that their budget submission or funding request is for a recovery task identified in an approved recovery plan, and is therefore part of a coordinated recovery effort to recover bull trout. In addition, Section 7 (a)(1) of the Endangered Species Act directs all Federal Agencies to use their authorities to further the purposes of the Act by implementing programs for the conservation of threatened or endangered species.

The following are definitions to column headings in the Implementation Schedule:

Priority Number: All priority 1 tasks are listed first, followed by priority 2 and priority 3 tasks.

Priority 1: All actions that must be taken to prevent extinction or to prevent the species from declining irreversibly in the foreseeable future.

Priority 2: All actions that must be taken to prevent a significant decline in species population, habitat quality, or some other significant negative effect short of extinction.

Priority 3: All other actions necessary to provide for full recovery (or reclassification) of the species.

Task Number and Task Description: Recovery tasks are numbered as in the recovery outline. Refer to the action narrative for task descriptions.

Task Duration: Expected number of years to complete the corresponding task. Study designs can incorporate more than one task, which when combined, may reduce the time needed for task completion.

Responsible or Participating Party: Federal, State, Native American Tribes, non-governmental organizations, or universities with responsibility or capability to fund, authorize or carry out the corresponding recovery task. Lead agencies are indicated in **bold type**. Additional identified agencies or parties are considered cooperators in restoration efforts.

Identified parties include:

BLM	U.S. Bureau of Land Management
BPT	Burns Paiute Tribe
ID	irrigation districts
NGOs	Non-governmental organizations
NRCS	Natural Resources Conservation Service
ODA	Oregon Department of Agriculture
ODEQ	Oregon Department of Environmental Quality
ODFW	Oregon Department of Fish and Wildlife
ODOF	Oregon Department of Forestry
OSP	Oregon State Police
OWRD	Oregon Water Resources Department
RUT	Recovery Unit Team
TMDLWG	Total Maximum Daily Load working group
USBOR	U.S. Bureau of Reclamation
USEPA	U.S. Environmental Protection Agency
USFS	United States Forest Service
USFWS	U.S. Fish and Wildlife Service
USGS	U.S. Geologic Service
WC	Watershed Councils

Cost Estimates: Cost estimates are rough approximations and are provided only for general guidance. Total costs are estimated for the duration of the task and also itemized annually for the next five years.

An asterisk (*) in the total cost column indicates ongoing tasks that are currently being implemented as part of normal agency responsibilities under existing authorities. Because these tasks are not being done specifically or solely for bull trout conservation, they are not included in the cost estimates. Some of these efforts may be occurring at reduced funding levels and/or in only a small portion of the watershed.

**BLANK
PAGE**

Implementation schedule for the bull trout recovery plan: Malheur Recovery Unit										
Task Priority	Task Number	Task Description	Task Duration (years)	Responsible Parties	Cost Estimates (in \$1,000 units)					Comments
					Total Costs	Year 1	Year 2	Year 3	Year 4	
1	1.1.1	Assess sediment sources in Malheur Basin affecting bull trout	10	USFS	30		10	10	10	Covered under existing programs
1	1.1.2	Stabilize roads, crossings, and other sources of sediment delivery	10		200		20	20	20	
1	1.2.1	Identify and implement opportunities for two-way passage at major dams	20	BOR, ID, ODFW, USFWS	50	50				Cost is for proposed study only; implementation costs unknown.
1	1.2.2	Provide passage at transportation/road-related barriers identified in completed and ongoing surveys	25	USFS, ODOT, Counties	50	50				Ongoing. Cost are for completion of surveys; project costs unknown at this time
1	1.2.3	Install appropriate fish screens and passage structures around diversions and/or remove related migration barriers	10	ODFW, BPT, USFS, landowners	100					Discussions ongoing. Cost estimate is for design only; project costs unknown at this time
1	1.3.4	Improve instream habitat	10	USFS, USFWS	2500	25	25	25	25	Ongoing
1	1.5.1	Assess current and historical effects of upland management on changes to the hydrograph, (e.g., timing and magnitude of peak flows) in all spawning tributaries	1	USFS, USFWS	50		50			

Implementation schedule for the bull trout recovery plan: Malheur Recovery Unit										
Task Priority	Task Number	Task Description	Task Duration (years)	Responsible Parties	Cost Estimates (in \$1,000 units)					Comments
					Total Costs	Year 1	Year 2	Year 3	Year 4	
1	2.5.1	Implement brook trout removal effort(s) wherever feasible and biologically supportable	25	ODFW, BPT	1000		40	40	40	
1	3.1.1	Incorporate bull trout recovery actions into The Oregon Plan for Salmon and Watersheds and the Pacific Northwest Power Planning Council Subbasin plans	25	ODFW	*					Currently providing monthly reports to Oregon Plan staff
1	3.3.1	Determine site-specific levels of competition and hybridization with introduced sport fish and assess impacts of those interactions	2	ODFW, USFWS	350		125	125		
1	4.1.1	Develop genetic management plan	3	USFWS, ODFW, BPT	300	100	100	100		
1	5.2.1	Identify site-specific threats that may be limiting bull trout in watersheds with historical bull trout habitat	3	USFS, RUT, USFWS	100					
1	5.2.3	Investigate potential for restoring historic prey base by reintroducing anadromous species	2-10	ODFW, BPT	30		30			Cost is for feasibility analysis only

Implementation schedule for the bull trout recovery plan: Malheur Recovery Unit										
Task Priority	Task Number	Task Description	Task Duration (years)	Responsible Parties	Cost Estimates (in \$1,000 units)					Comments
					Total Costs	Year 1	Year 2	Year 3	Year 4	
1	5.5.2	Conduct regular surveys in potential habitat in the Malheur Core Area where bull trout status is unknown or recolonization is anticipated	25	RUT, USFWS, USFS, BLM	200					Periodic re-surveys during life of recovery plan
1	5.5.4	Determine consequences of genetic fragmentation/isolation due to human-made barriers	5	USFWS, ODFW	500		100	100	100	
1	6.1.4	Identify and pursue opportunities to implement recovery strategies	25	USFWS, RUT	*					
1	7.1.1	Develop an annual work plan to support implementation in the Malheur Recovery Unit	5	RUT	*					
2	1.1.4	Increase monitoring of sediment inputs on the Malheur National Forest	20	USFS	200	10	10	10	10	
2	1.3.1	Restore shade and canopy, riparian cover, and native vegetation in all bull trout spawning areas	10	USFS, BLM, BPT, WC, landowners	250	25	25	25	25	Efforts currently underway on public and tribal lands
2	1.3.2	Reduce grazing impacts in all bull trout spawning areas	5	USFS, BPT, landowners	500		100	100	100	

Implementation schedule for the bull trout recovery plan: Malheur Recovery Unit										
Task Priority	Task Number	Task Description	Task Duration (years)	Responsible Parties	Cost Estimates (in \$1,000 units)					Comments
					Total Costs	Year 1	Year 2	Year 3	Year 4	
2	1.3.3	Review habitat information to identify and prioritize opportunities for channel restoration in Lake Creek	1	USFS, ODFW, BPT, USFWS	30		30			Cost for study and design only; implementation costs unknown at this time
2	1.4.1	Review reservoir operational concerns and provide operating recommendations through Federal consultation or other means	25	BOR, USFWS	*					Studies underway
2	1.4.2	Establish/provide instream flows downstream from reservoirs and stabilize flow regimes	25	USBR, ODFW	100					
2	2.3.1	Develop and implement an educational effort to address problems and consequences of unauthorized fish introductions	1	USFWS, ODFW	10		10			
2	3.2.1	Maintain bull trout as high priority for Oregon's Cooperative Enforcement Program	25	ODFW, OSP	*					Priorities set at annual coordination meetings
2	5.5.3	Determine life history requirements of resident and migratory bull trout local populations in the Malheur Core Area	5	ODFW, BPT, USFWS	250		50	50	50	

Implementation schedule for the bull trout recovery plan: Malheur Recovery Unit										
Task Priority	Task Number	Task Description	Task Duration (years)	Responsible Parties	Cost Estimates (in \$1,000 units)					Comments
					Total Costs	Year 1	Year 2	Year 3	Year 4	
2	6.1.1	Provide long-term habitat protection	25	ODFW, BPT, USFS, BLM, USFWS	1,250		50	50	50	
2	6.1.3	Develop educational materials on bull trout and their habitat needs to provide to landowners and interested public parties	2	ODFW, USFS, USFWS	40		20	20		
2	6.1.5	Integrate watershed analyses and assessments and restoration activities on public and private lands	25	RUT, DEQ, WC	*					
3	1.1.3	Assess and mitigate effects on bull trout from non-point source pollution	10	DEQ, WC, ODA, NRCS, USFWS	200	20	20	20	20	Covered under TMDL and SB1010 plans
3	1.2.4	Restore connectivity and opportunities for migration by improving instream flows and/or water rights	25	ODFW, OWRD, BPT	1000	40	40	40	40	Ongoing
3	3.1.2	Coordinate bull trout recovery with recovery efforts, management plans, etc. of other species such as redband trout	25	ODFW, USFWS	*					Covered under existing management plan.

Implementation schedule for the bull trout recovery plan: Malheur Recovery Unit										
Task Priority	Task Number	Task Description	Task Duration (years)	Responsible Parties	Cost Estimates (in \$1,000 units)					Comments
					Total Costs	Year 1	Year 2	Year 3	Year 4	
3	3.2.2	Increase information outreach to anglers	5	ODFW, USFS, BLM, USFWS	50		50			Costs are for development of materials; implementation would be an ongoing process
3	3.2.3	Improve and implement fisheries management guidelines and policies designed to protect native species	25	ODFW, USFWS, USFS, BLM	*					
3	5.1.1	Evaluate effectiveness of different habitat restoration techniques in restoring channel functions and local bull trout populations in the Malheur Core Area	10	USFS, BLM, BPT, USFWS	200		20	20	20	
3	5.2.2	Determine the movement and seasonality of use of different habitat types by adult and sub-adult migratory bull trout in multiple streams, with emphasis on reservoirs and mainstem rivers	3	RUT, ODFW, USFW, USFS, BLM	*					In progress
3	5.5.1	Review and update databases for bull trout distribution records	25	RUT	250		5	5	5	Basic database has been established and incorporated into StreamNet (2002)

Implementation schedule for the bull trout recovery plan: Malheur Recovery Unit										
Task Priority	Task Number	Task Description	Task Duration (years)	Responsible Parties	Cost Estimates (in \$1,000 units)					Comments
					Total Costs	Year 1	Year 2	Year 3	Year 4	
3	5.5.5	Evaluate food web interactions in drainages most affected by introduced fishes, reservoir operations, loss of anadromous species (prey base/nutrients), etc	25	BOR, USGS	*					Ongoing studies
3	6.1.2	Work cooperatively with the Burns-Paiute tribal government to implement recovery actions	25	RUT, USFWS	*					
3	7.3.1	Periodically review progress towards recovery goals and assess recovery task	25	USFWS	*					

REFERENCES

- Bartholomew, J. 2001. Fish health risk assessment - Oregon State University 2000 progress report. *In: Portland General Electric Pelton Round Butte Fisheries Workshop*, Spring 2001. Portland General Electric Company. Portland, Oregon.
- (BLM) U.S. Bureau of Land Management. 1995. Riparian Area Management: Process for Assessing Proper Functioning Condition. Report Number T-1737-09.
- (BLM) U.S. Bureau of Land Management. 2002. Final report on 2001 grazing management in North Fork Malheur River allotments. Vale, Oregon.
- Bowers, W., P. Dupee', M.L. Hanson, and R. Perkins. 1993. Bull trout population summary Malheur River Basin. Unpublished report. Oregon Department of Fish and Wildlife, Hines, Oregon.
- Buchanan, D.V., M.L. Hanson, and R.M. Hooton. 1997. Status of Oregon's Bull Trout. Oregon Department of Fish and Wildlife, Portland, Oregon.
- Buckman, R. C., W. E. Hosford, and P.A. Dupee. 1992. Malheur River bull trout investigations. Pages 45 - 57 *In: Howell, P. J., and D.V. Buchanan, editors. Proceedings of the Gearhart Mountain bull trout workshop. Oregon Chapter of the American Fisheries Society, Corvallis.*
- California Department of Fish and Game. 1985. Draft programmatic environmental impact report. Rotenone use for fisheries management. Pesticides Investigations Unit, Rancho Cordova, California.
- Crow, J.F., and M. Kimura. 1970. An introduction to population genetics theory. Harper and Row, New York.
- Franklin, I.R. 1980. Evolutionary changes in small populations. Pages 135-149 *In: M.E. Soule, and B.A. Wilcox, editors. Conservation Biology: An Evolutionary-Ecological perspective. Sinauer Associates, Sunderland, Massachusetts.*
- Franklin, J.F., and C.T. Dyrness. 1984. Natural Vegetation of Oregon and Washington. Oregon State University Press. Corvallis, Oregon.

- Gunckel, S.L. 2000. Feeding behavior and diet of native bull trout *Salvelinus confluentus* and introduced brook trout *S. fontinalis* in two eastern Oregon streams. Master of Science dissertation. Oregon State University, Corvallis.
- Hanson, M.L., R.C. Buckman, and W.E. Hosford. 1990. Malheur River basin fish management plan. Oregon Department of Fish and Wildlife. Hines, Oregon.
- Kostow, K. 1995. Biennial report on the status of wild fish in Oregon. Oregon Department of Fish and Wildlife. Portland, Oregon.
- Lande, R. 1988. Genetics and demography in biological conservation. *Science* 241:1455-1460.
- Malheur-Owyhee Watershed Council. 1999. Malheur Basin Action Plan. Ontario, Oregon.
- Meffe, G.K., and C.R. Carroll. 1994. Principles of conservation biology. Sinauer Associates, Inc. Sunderland, Massachusetts.
- Northwest Power Planning Council (NWPPC). 2002. Draft Malheur Subbasin Summary. Portland, Oregon.
- Oregon Department of Agriculture (ODA). 2001. Malheur River basin agricultural water quality management area plan. Developed with assistance from Malheur River Basin Agricultural Water Quality Local Advisory Committee and Malheur County Soil and Water Conservation District. Portland, Oregon.
- Perkins, R. 2000. Malheur River bull trout limiting factors analysis. Oregon Department of Fish and Wildlife. Ontario, Oregon.
- Ratliff, D.E., and P.J. Howell. 1992. The status of bull trout populations in Oregon. Pages 10 - 17 *In*: Howell, P.J., and D.V. Buchanan, editors. Proceedings of the Gearhart Mountain bull trout workshop. Oregon Chapter of the American Fisheries Society, Corvallis.
- Rieman, B.E., and Allendorf, F.W. 2001. Effective population size and genetic conservation criteria for bull trout. *North American Journal of Fisheries Management* 21:756-764.
- Rieman, B.E., and J.D. McIntyre. 1993. Demographic and habitat requirements for conservation of bull trout. General Technical Report INT-302. U.S. Forest Service. Intermountain Research Station, Boise, Idaho.
- Schwabe, L., M. Tiley, and R. Perkins. 2000. Malheur River Basin Cooperative Bull Trout/Redband Trout Research Project, FY1999 Annual Report. Report to Bonneville

Power Administration, Portland, Oregon. Contract No. 00006313, Project No. 199701900, 120 electronic pages (BPA Report DOE/BP-00006313-1).

Schwabe, L., S. Namitz, J. Fenton, R. Perkins, P. Spruell, and A. Maxwell. 2001. Evaluate the life history of native salmonids within the Malheur Subbasin. FY 2000 Annual Report, Report to Bonneville Power Administration, Portland, Oregon. Contract No. 00006313, Project No. 9701900/9701901).

Soule, M.E. 1980. Thresholds for survival: maintaining fitness and evolutionary potential. Pages 151-170 *In*: M. E. Soule, and B. A. Wilcox, editors. Conservation biology: An evolutionary-ecological perspective. Sinauer and Associates, Sunderland, Massachusetts

Spruell, P., and F.W. Allendorf. 1997. Nuclear DNA analysis of Oregon bull trout. Final report to Oregon Department of Fish and Wildlife. Division of Biological Sciences, University of Montana, Missoula.

Spruell, P., A.R. Hemmingsen, P.J. Howell, N. Kanda, and F.W. Allendorf. 2002. Conservation genetics of bull trout: geographic distribution of variation at microsatellite loci. Conservation Genetics (in press).

State of Oregon (Nicholas, J.W., Principal Writer). 1997. The Oregon plan (Oregon Coastal Salmon Restoration Initiative). Oregon Governor's Office, Salem, OR. 3,000+ pp. Also available at the following web site: <http://www.oregon-plan.org/>.

StreamNet. 2002. Fish data for the Northwest. <http://www.streamnet.org/>. Accessed August 28, 2002. Administered by the Pacific States Marine Fisheries Commission. Gladstone, Oregon.

Tinneswood, B., and R. Perkins. 2001. Bull trout spawning survey report, 2000. Unpublished report. Oregon Department of Fish and Wildlife. Ontario, Oregon.

(USBOR) U.S. Bureau of Reclamation. 1998. Bureau of Reclamation operations and maintenance in the Snake River Basin above Lower Granite Reservoir biological assessment. Boise, Idaho.

(USBOR) U.S. Bureau of Reclamation. 2001. Structural alternatives for establishment of a conservation pool for bull trout at Beulah Reservoir in the Malheur River Basin. Appraisal Report. Boise, Idaho.

(USFS) U.S. Forest Service. 1995. Inland native fish strategy. Environmental assessment. Intermountain, Northern, and Pacific Northwest Regions.

- (USFS) U.S. Forest Service. 1999a. Bull trout biological assessment. Malheur River ongoing projects. Malheur National Forest, Prairie City Ranger District. Prairie City, Oregon.
- (USFS) U.S. Forest Service. 1999b. Bull trout biological assessment. North Fork Malheur River ongoing projects. Malheur National Forest, Prairie City Ranger District. Prairie City, Oregon.
- (USFS) U.S. Forest Service. 2000. Malheur headwaters watershed analysis. Prairie City Ranger District, Malheur National Forest. Prairie City, Oregon.
- (USFS) U.S. Forest Service. 2002. Biological assessment for merit vegetation project. Prairie City Ranger District, Malheur National Forest. Prairie City, Oregon.
- (USFWS) U.S. Fish and Wildlife Service. 1999. Biological Opinion on the continued operation and maintenance of the Bureau of Reclamation facilities in the Snake River Basin upstream of the Corps of Engineers Lower Granite Reservoir, (File #1009.2700) (1-4-98-F-1) Final Document. Snake River Basin Office, Boise, Idaho.
- (USFWS) U.S. Fish and Wildlife Service. 2001. Biological Opinion for effects of Malheur National Forest's 2001 grazing management program for bull trout. (OLS Number: 01-3912) Oregon Fish and Wildlife Office, Portland, Oregon.

Literature

Burns Paiute Tribe. No date. Paiute wadatika. The history and cultural background of the Burns Paiute Tribe. 20 pages.

Rieber, R., U.S. Bureau of Reclamation. 2000. Comments on Malheur Chapter working draft to Mary Hanson. 2 pages.

Young, D., U.S. Fish and Wildlife Service. 2000. Comments on Malheur Chapter working draft to Mary Hanson.

Personal Communications

Bowers, W., Oregon Department of Fish and Wildlife. 2002. Portland, Oregon.

Butcher, D., Oregon Department of Environmental Quality. 2002. Portland, Oregon.

Miller, A., U.S. Forest Service. 2002. Subject: bull trout surveys. Prairie City, Oregon.

Miller, A., U.S. Forest Service. 2002. Subject: bull trout size classes observed. Prairie City, Oregon

Perkins, R., Oregon Department of Fish and Wildlife. 2002. Ontario, Oregon.

Rieber, R., U.S. Bureau of Reclamation. 2002. Boise, Idaho.

Schwabe, L., Burns Paiute Tribe. 2002. Burns, Oregon.

**APPENDIX A: Historical and current fish species found in the Malheur River Basin
excerpted from NWPPC 2001.**

Common Name	Scientific Name	Special Status	Distribution
Lampreys Family Petromyzontidae			
Pacific Lamprey	<i>Lampetra tridentata</i>	Extirpated	
Minnows Family Cyprinidae			
Chiselmouth	<i>Acrocheilus alutaceus</i>		Lower Malheur River
Northern Pikeminnow	<i>Ptychocheilus oregonensis</i>		Lower sections of major subbasins
Common Carp ²	<i>Cyprinus carpio</i>		Lower Malheur River
Long-nosed Dace	<i>Rhinichthys cataractae</i>		Lower sections of major subbasins
Speckled Dace	<i>Rhinichthys osculus</i>		Lower sections of major subbasins
Redside Shiner	<i>Richardsonius balteatus</i> <i>balteatus</i>		Lower sections of major subbasins
Loaches Family Cobitidae			
Oriental Weatherfish ¹	<i>Misgurnus anguillicaudatus</i>		Irrigation and drain ditches in lower Subbasin
Suckers Family Catostomidae			
Largescale Sucker	<i>Catostomus macrocheilus</i>		Larger river and reservoirs
Largescale Sucker	<i>Catostomus macrocheilus</i>		Larger river and reservoir
Bridgelip Sucker	<i>Catostomus columbianus</i>		Lower sections of major subbasins
Bullhead catfishes Family Ictaluridae			
Channel Catfish ¹	<i>Ictalurus punctatus</i>		Warm Springs Reservoir, Bully Creek Reservoir, and lower Malheur River
Brown Bullhead ¹	<i>Ameiurus nebulosus</i>		Warm Springs and Bully Creek Reservoirs
Trouts Family Salmonidae			
Chinook Salmon	<i>Oncorhynchus tshawytscha</i>	Extirpated	

APPENDIX A: Historical and current fish species found in the Malheur River Basin excerpted from NWPPC 2001.

Common Name	Scientific Name	Special Status	Distribution
Coho Salmon	<i>Oncorhynchus kisutch</i>	Extirpated	
Steelhead	<i>Oncorhynchus mykiss</i>	Extirpated	
Columbia River Redband Trout	<i>Oncorhynchus mykiss</i>	State Sensitive	Higher elevation areas of most major subbasins
Hatchery Rainbow Trout ¹	<i>Oncorhynchus mykiss</i>		Malheur, Pole Creek, Beulah, Warm Springs, Murphy, Cottonwood Reservoirs, 9 small BLM stock ponds, and Malheur River from Gold Creek to Warm Springs Dam
Bull Trout	<i>Salvelinus confluentus</i>	Federal Threatened	Headwaters of North Fork and Logan Valley streams
Brook Trout ¹	<i>Salvelinus fontinalis</i>		Logan Valley streams
Whitefish	<i>Prosopium williamsoni</i>		Lower sections of North Fork, Upper Malheur, and lower Malheur River
Sculpin	Family Cottidae		
Shorthead Sculpin	<i>Cottus confusus</i>		Headwater areas of perennial streams
Mottled Sculpin Nongame	<i>Cottus bairdi</i>		Headwater areas of perennial streams
Bass	Family Cyprinodontidae		
Largemouth Bass ¹	<i>Micropterus salmoides</i>		Warm Springs and Bully Creek Reservoirs
Smallmouth ¹ Bass	<i>Micropterus dolomieu</i>		Warm Springs and Bully Creek Reservoirs
White Crappie ¹	<i>Pomoxis annularis</i>		Warm Springs and Bully Creek Reservoirs
Bluegill ¹	<i>Lepomis macrochirus</i>		Warm Springs and Bully Creek Reservoirs
Perch	Family Percidae		
Yellow Perch ¹	<i>Perca flavescens</i>		Warm Springs and Bully Creek Reservoirs

¹ Introduced species

APPENDIX B: List of barriers in the Malheur Recovery Unit known or suspected to impact bull trout. (Agency and Warm Springs Dams and the Drewsey diversion are priorities.)

Subbasin	Location	Type of Barrier	Comment
North Fork Malheur River	Agency Dam	Storage Dam	
	North Fork Malheur	Diversions	
Upper Malheur River	Warm Springs Dam	Storage Dam	
	Drewsey Diversion	Diversion Dam	
	Summit Creek	Road crossings, thermal barrier near mouth	
	Bosnberg Creek	Railroad grade crossing	
	Lake Creek	Thermal barrier near mouth	
	McCoy Creek	Thermal barrier near mouth	
	Malheur River	Diversions	

APPENDIX C: List of Instream Water Rights established to benefit bull trout (BuT) and redband trout (RT).					
Application #	Certificate #	Species	Stream	From	To
71456	73348	BuT,RT	N FK MALHEUR R > MALHEUR R	L MALHEUR R	BEULAH RES
70304	73338	BuT,RT	N FK MALHEUR R > MALHEUR R	CRANE CR	L MALHEUR R
70306	73340	BuT,RT	CRANE CR > N FK MALHEUR R	L CRANE CR	MOUTH
70307	73341	BuT,RT	LITTLE CRANE CR > CRANE CR	HEADWATERS	MOUTH
70308	73342	BuT,RT	ELK CR > N FK MALHEUR R	HEADWATERS	MOUTH
70309	73343	BuT,RT	SWAMP CR > N FK MALHEUR R	HEADWATERS	MOUTH
70349	68360	BuT,RT	MALHEUR R (M FK) > MALHEUR R	LOGAN VALLEY	UP DREWSEY DIV
70352	68363	BuT,RT	BIG CR > LAKE CR	HEADWATERS	USFS RD 16 XING
70303	73337	BuT,RT	N FK MALHEUR R > MALHEUR R	USFS RD 16 XING	CRANE CR
70351	68362	BuT,RT	LAKE CR > MALHEUR R	HEADWATERS	USFS RD 16 XING

APPENDIX D: Oregon Department of Fish and Wildlife instream water right priorities for bull trout (BuT) and redband trout (RT).**North Fork Malheur River**

Stream	Begin	End	Priority	Species	Spawning	Incubation	Juvenile Rearing	Subadult Rearing	Adult Rearing
Sheep Creek	mouth	headwaters	1	BuT	yes	yes	yes	probably	yes
N. F. Malheur R.	FS Rd 16 bridge	FS Rd 1370 bridge	1	BuT	yes	yes	yes	probably	yes
Horseshoe Creek	mouth	headwaters	1	BuT	yes	yes	yes	probably	yes
Swamp Creek Trib	mouth	headwaters	1	BuT	unknown	unknown	yes	probably	yes
Flat Creek	mouth	headwaters	1	BuT	unknown	unknown	yes	probably	yes
Crane Creek	mouth	headwaters	2	BuT	unknown	unknown	yes	probably	probably
Cow Creek	mouth	headwaters	3	RT			probably	probably	yes
Little Cow Creek	mouth	headwaters	3	RT					
Spring Creek	mouth	headwaters	3	RT					
Buttermilk Creek	mouth	headwaters	3	RT					
Bear Creek	mouth	headwaters	3	RT					
Little Malheur River									
Lost Creek	mouth	headwaters	3	RT					
Bridge Creek	mouth	headwaters	3	RT					
Squaw Creek	mouth	headwaters	3	RT					
Rock Creek	mouth	headwaters	3	RT					
South Bullrun Creek	mouth	headwaters	3	RT					
Lunch Creek	mouth	headwaters	3	RT					
Larch Creek	mouth	headwaters	3	RT					
Canteen Creek	mouth	headwaters	3	RT					

APPENDIX D: Oregon Department of Fish and Wildlife instream water right priorities for bull trout (BuT) and redband trout (RT)

Camp Creek	mouth	headwaters	3	RT					
Upper Malheur River									
Meadow Fork	mouth	headwaters	1	BuT	yes	yes	yes	probably	yes
Snowshoe Creek	mouth	headwaters	1	BuT	yes	yes	yes	probably	yes
Crooked Creek	mouth	headwaters	2	BuT	unknown	unknown	probably	probably	unknown
Lake Creek	mouth	FS Rd 16 Road	2	BuT	unknown	unknown	unknown	maybe	probably
Big Creek	mouth	FS Rd 16 Road	2	BuT	unknown	unknown	maybe	probably	probably
Corral Basin Creek	mouth	headwaters	3	BuT	unknown	unknown	unknown	unknown	unknown

APPENDIX E: List of current and potential (*) local bull trout populations in the Malheur Recovery Unit. Bull trout in the Malheur Recovery Unit exhibit a fluvial life history.

Subbasin	Streams	Habitat Use
North Fork Malheur	Mainstem North Fork, mouth to Agency Dam	Migration/overwintering/foraging
	Beulah Reservoir	Migration/overwintering/foraging
	Mainstem North Fork, Beulah Reservoir to confluence with Crane Creek	Migration/overwintering/foraging
	Mainstem North Fork upstream of confluence with Crane Creek	Spawning/rearing
	Cow Creek	Rearing
	Horshoe Creek	Spawning/rearing
	Flat Creek	Spawning/rearing
	Swamp Creek	Spawning/rearing
	Sheep Creek	Spawning/rearing
	Elk Creek and tributaries North Fork and South Fork Elk Creeks	Spawning/rearing
	Crane Creek	Migration/foraging
	Little Crane Creek	Spawning/rearing
Upper Malheur River	Little Malheur*	Migration/overwintering/foraging
	Mainstem Malheur above Drewsey	Migration/overwintering/foraging
	Mainstem* Malheur from Drewsey to Namorf Dam	Migration/overwintering/foraging
	Warm Springs Reservoir*	Overwintering/foraging

APPENDIX E: List of current and potential (*) local bull trout populations in the Malheur Recovery Unit. Bull trout in the Malheur Recovery Unit exhibit a fluvial life history.

Subbasin	Streams	Habitat Use
	Lake Creek	Spawning/rearing
	Big Creek	Spawning/rearing
	Meadow Fork Big Creek	Spawning/rearing
	Snowshoe Creek (Big Creek tributary)	Spawning/rearing
	Summit Creek	Spawning/rearing
	Crooked Creek*	Spawning/rearing
	Bosonberg Creek*	Spawning/rearing
	McCoy Creek*	Spawning/rearing
	Corral Basin* (Tributary to Big Creek)	Spawning/rearing

APPENDIX F: List of Chapters

- Chapter 1 - Introductory
- Chapter 2 - Klamath Recovery Unit, Oregon
- Chapter 3 - Clark Fork Recovery Unit, Montana, Idaho, and Washington
- Chapter 4 - Kootenai Recovery Unit, Montana and Idaho
- Chapter 5 - Willamette Recovery Unit, Oregon
- Chapter 6 - Hood Recovery Unit, Oregon
- Chapter 7 - Deschutes Recovery Unit, Oregon
- Chapter 8 - Odell Lake Recovery Unit, Oregon
- Chapter 9 - John Day Recovery Unit, Oregon
- Chapter 10 - Umatilla-Walla Walla Rivers Recovery Unit, Oregon and Washington
- Chapter 11 - Grande Ronde Recovery Unit, Oregon
- Chapter 12 - Imnaha-Snake Rivers Recovery Unit, Oregon
- Chapter 13 - Hells Canyon Complex Recovery Unit, Oregon and Idaho
- Chapter 14 - Malheur Recovery Unit, Oregon**
- Chapter 15 - Coeur d'Alene Recovery Unit, Idaho
- Chapter 16 - Clearwater Recovery Unit, Idaho
- Chapter 17 - Salmon Recovery Unit, Idaho
- Chapter 18 - Southwest Idaho Recovery Unit, Idaho
- Chapter 19 - Little Lost Recovery Unit, Idaho
- Chapter 20 - Lower Columbia Recovery Unit, Washington
- Chapter 21 - Middle Columbia Recovery Unit, Washington
- Chapter 22 - Upper Columbia Recovery Unit, Washington
- Chapter 23 - Northeast Washington Recovery Unit, Washington
- Chapter 24 - Snake River Washington Recovery Unit, Washington
- Chapter 25 - Saint Mary - Belly Recovery Unit, Montana

END

04-10-03